


Researchers develop new method to diagnose, monitor chronic wounds

November 20 2018, by Matt McGowan



Kyle Quinn in his lab at the University of Arkansas. Credit: University of Arkansas

Researchers at the University of Arkansas have developed a new approach to diagnosing and monitoring chronic skin wounds, such as

diabetic foot ulcers and pressure wounds.


Chronic skin wounds affect more than 150 million people worldwide and cost approximately \$25 billion in health care annually in the United States alone. These non-healing wounds are characterized by inflamed [tissue](#), poor blood circulation, callus formation or infection.

Current clinical approaches for diagnosing and monitoring these wounds do not provide critical diagnostic information about how they develop or why they do not heal.

Kyle Quinn, assistant professor of biomedical engineering, and Jake Jones, a doctoral student in biomedical engineering, have identified how cell metabolism changes over time during the healing process of wounds, by using a powerful imaging system capable of measuring changes in cellular metabolism.

Quinn and Jones used a process called label-free multiphoton microscopy to view tissue in three dimensions at the cellular level and generate 3D maps of wound metabolism. Label-free means the tissue was not artificially labeled with a fluorescent molecule. Multiphoton microscopy can facilitate measurements of naturally occurring cell and tissue autofluorescence, which is the emission of light by molecules naturally present in [cells](#) and tissue when those molecules have absorbed light.

In this study, autofluorescence imaging of two molecules – [nicotinamide adenine dinucleotide](#) (NADH) and flavin adenine dinucleotide (FAD) – were used to monitor cell metabolism. This was accomplished through a measurement called an optical redox ratio, which quantifies the relative concentrations of NADH and FAD. The researchers monitored wound metabolism in mice over a period of 10 days.



Credit: University of Arkansas

"To our knowledge, this is the first study showing changes in the optical redox ratio of tissue within live animals imaged through repeated sessions," Quinn said. "We consider this a critical step in demonstrating the diagnostic potential of multiphoton microscopy for clinical wound care."

Complications associated with diabetes are a common cause of chronic [wounds](#). For this reason, Quinn and Jones's study included diabetic mice, in addition to healthy, control mice.

The researchers observed differences between diabetic and control mice during the rebuilding of wound tissue. Changes in the optical redox ratio and NADH fluorescence among the diabetic mice indicated that cells remained at the wound edge, growing and dividing, rather than migrating over the wound to restore the skin's protective barrier function.

"The ability of multiphoton microscopy to non-invasively collect structural and metabolic data suggests that it might have broader applications for wound care and dermatology," said Quinn. "Providing quantitative metrics of cell function and wound composition has potential to supplement current standard of care and guide treatment

options."

The study was published in *Communications Biology*.

More information: Jake D. Jones et al. In vivo multiphoton microscopy detects longitudinal metabolic changes associated with delayed skin wound healing, *Communications Biology* (2018). [DOI: 10.1038/s42003-018-0206-4](https://doi.org/10.1038/s42003-018-0206-4)

Provided by University of Arkansas

Citation: Researchers develop new method to diagnose, monitor chronic wounds (2018, November 20) retrieved 20 April 2024 from <https://medicalxpress.com/news/2018-11-method-chronic-wounds.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.