


# **Precision neuroengineering enables reproduction of complex brain-like functions in vitro**

November 14 2018

---

## Re-creation of *neuronal circuits* in vitro



**T**he brain is a modular network, which can operate as a localized circuit (segregation) or in coordination with other circuits (integration). Researchers have reproduced this behaviour by growing neurons on an adhesive molecule pattern divided into four modules connected by tracks.

The seeded neurons spontaneously activate and transfer information through the existing connections between them and the different modules, reproducing in vitro the complex balance of integration and segregation observed in the brain.

Researchers have reproduced a brain-like behaviour by growing neurons on a pattern divided into four modules connected by tracks. Credit: Universitat de Barcelona / Tohoku University.

One of the most important and surprising traits of the brain is its ability to dynamically reconfigure the connections to process and respond properly to stimuli. Researchers from Tohoku University (Sendai, Japan) and the University of Barcelona, using neuroengineering tools, have created in vitro neural circuits that reproduce the capacity of segregation and integration of brain circuits and which allow researchers to understand the keys of dynamic reconfiguration. The study has been published in *Science Advances*.

Dynamic reconfiguration is understood as the strengthening or weakening of connections by increasing or decreasing neuronal activity. In the [brain](#), an increase in the cohesion of [neuronal circuits](#) is known as integration, and a decrease is known as segregation. "This study shows the importance of modular organization to maximize the flexibility of a neural circuit. It also shows the potential of in vitro tools and biophysical models to progress in the understanding of collective phenomena in such a fascinating and rich complex system like the brain" says Jordi Soriano, researcher at the Institute of Complex Systems of the UB (UBICS) and co-author of the study.

Integration is linked to the rapid exchange of information between different and distant circuits, while segregation is linked to information processing in localized [circuits](#). The brain goes from a segregated state to an integrated one depending on the nature and strength of [stimuli](#). Dynamic reconfiguration avoids creating and destroying physical connections, a strategy as ineffective as energetically costly. Therefore, stimuli that arrive through sight, hearing and smell, are processed in a segregated way in the cerebral cortex to partially or totally integrate according to the needs. While we watch a film, we integrate images and sounds, ignoring smells and other stimuli. However, when we notice a smell of something burning the brain enters a warning state so that we integrate and analyse all the information available to make fast decisions.

Despite the importance of integration and segregation, biophysical mechanisms linked to dynamic reconstructions are still not well understood. Also, another unknown element is the sensitivity of integration-segregation ability regarding the amount of existing physical connections between brain regions.

The in vitro brain model the researchers developed consists of four interconnected modules. Each module represents a specialized neuronal circuit (sight or hearing, for instance). These four modules are covered by adhesive proteins and nutrients where neurons develop. These neurons are connected among them inside a module and with other neurons in other modules. Precision neuroengineering allows controlling the connections that go from one module to another, and therefore, it enables adjusting the level of physical coupling between modules. The stimuli in this model correspond to neural spontaneous activations.

Using fluorescence calcium imaging to detect neural activations, researchers studied the ability of the circuit to integrate or segregate spontaneously depending on the level of connectivity between modules and other factors. "What we saw is that the circuit is fully integrated or segregated when the number of connections between modules is too big or too small. The optimal circuit is that where four modules have a connectivity just below the minimum to integrate, so that neural activity pulses are enough to reinforce connections and complete integration. In effect, this optimal circuit—spontaneously activated—works in a regime where [integration](#) and [segregation](#) coexist. Of course, the observed dynamics are still far from the complexity in the actual brain, but we were able to gain insight into the fundamental mechanisms shaping brain dynamics", notes Hideaki Yamamoto, researcher at Tohoku University.

**More information:** H. Yamamoto, S. Moriya, K. Ide, T. Hayakawa, H. Akima, S. Sato, S. Kubota, T. Tanii, M. Niwano, S. Teller, J. Soriano, A. Hirano-Iwata. "Impact of modular organization on dynamical richness in

cortical networks". *Science Advances* DOI: 10.1126/sciadv.aau4914 ,  
[advances.sciencemag.org/content/4/11/eaau4914](https://advances.sciencemag.org/content/4/11/eaau4914)

Provided by University of Barcelona

Citation: Precision neuroengineering enables reproduction of complex brain-like functions in vitro (2018, November 14) retrieved 25 April 2024 from  
<https://medicalxpress.com/news/2018-11-precision-neuroengineering-enables-reproduction-complex.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.