

PTSD common among those who suffer tear in the aorta's wall

November 13 2018



The sharp and sudden pain from an aortic dissection, along with the

emergency treatment that follows, can lead to post-traumatic stress disorder years later, a new study finds.

An [aortic dissection](#) is a life-threatening condition in which a tear in the wall of the aorta—the major artery carrying blood out of the [heart](#)—allows blood to rush between the wall's layers. Most people who have one say they feel a sudden ripping or stabbing chest pain that sometimes spreads to the back. It is usually treated with surgery and medication.

In a study of aortic dissection patients treated at the Cleveland Clinic, researchers reached out to 295 people with an online survey that included four questions commonly used to screen for PTSD.

Patients were asked whether they startled easily, or felt constantly on guard since their dissection. They also were asked about feeling numb or detached from others, whether they had nightmares or unwelcome thoughts about their dissection, or if they went out of their way to avoid situations that reminded them of it.

About 44 percent of those surveyed, or 129 people, answered all four screening questions. Of that group, 22 percent screened positive for PTSD. The results "absolutely shocked" researchers, said Selena Pasadyn, a medical student at the Cleveland Clinic Lerner College of Medicine and the study's lead investigator.

"We're advancing a lot in our care for these patients. We're able to get them through this acute event, but it's also our responsibility as physicians to think about their long-term quality of life," Pasadyn said. "This is a cry to physicians to consider PTSD, to screen for it, and to refer and treat appropriately."

She said she also hopes to raise awareness among patients so they can become better advocates for themselves.

"They need to be aware that they may feel watchful. They may feel on guard or startled," she said. "They need to know these are normal feelings, but they're also treatable feelings."

The average age of the patient responding to the survey was 54, and the average time that had passed since their aortic dissection was about seven years, she said.

The findings were presented Saturday at the American Heart Association's Scientific Sessions conference, in Chicago.

While the study is one of only a few to look specifically at aortic dissection patients, related research has found that about 15 to 25 percent of heart attack and stroke survivors also screen positive for PTSD, said Donald Edmondson, director of Columbia University's Center for Behavioral Cardiovascular Health in New York City.

An aortic dissection has all of the characteristics of a PTSD-inducing cardiovascular event, he said.

"Any type of event that is sudden onset, highly painful, and life-threatening has the potential to cause PTSD," said Edmondson, who was not involved in the new study. "But what's unique and special about medical events like this is that unlike combat or sexual assault, where people can find a safe place where they're not at risk, someone who's had an aortic dissection or a heart attack or stroke can never get away from their heart (or brain)."

Edmondson said that [heart attack](#) survivors with PTSD are at double the risk of having another attack or dying within the next year than a survivor who is not under similar distress.

There are two reasons for this, he said. One has to do with the body's

natural reaction to stress. Just thinking about a traumatic event can make some people hyperventilate or become lightheaded.

"Often when people become aware of their physiological signals, like feeling short of breath, it distresses them. It makes them worry they're going to have another cardiovascular event, which of course makes their heart rate go faster," Edmondson said. "That vicious cycle appears to contribute to secondary risk."

Heart attack and [stroke survivors](#) with PTSD also are less likely to take medicine prescribed to help prevent future events.

"Their medications serve as reminders of their [cardiovascular event](#) and the fact that they're still at risk," Edmondson said. "They try to avoid thinking about that, even though they think about it all the time."

Copyright is owned or held by the American Heart Association, Inc., and all rights are reserved. If you have questions or comments about this story, please email editor@heart.org.

Citation: PTSD common among those who suffer tear in the aorta's wall (2018, November 13) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2018-11-ptsd-common-aorta-wall.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|