

# Research reveals that 1 in 4 suicide attempts are associated with perceptual difficulties

November 28 2018

---



Credit: CC0 Public Domain

Researchers at RCSI (Royal College of Surgeons in Ireland) have revealed that one quarter of suicide attempts are associated with dysfunction in how the brain interprets basic perceptual information,

such as what we see, hear and think. The research shows that this dysfunction can predict suicidal behaviour, and offers new prospects for treatment and suicide prevention. The research has today been published in *JAMA Psychiatry*.

Symptoms such as depressed mood, feelings of worthlessness and a sense of hopelessness are well documented in suicidal behaviour. However, in an analysis of more than 80,000 people, new [research](#) has revealed that one quarter of individuals who attempted or died by [suicide](#) had problems in basic sensory [experiences](#), such as hearing or seeing things that aren't really there—otherwise known as 'perceptual abnormalities'.

Such episodes are not necessarily associated with [psychotic illnesses](#) or depression and can occur in people who do not experience mental illness, according to the study.

Research over the past 15 years has shown that experiences such as 'hearing voices' are far more common than previously thought—about 5-7% of the general population report at least occasionally having experiences such as hearing voices. For some people, these experiences emerge when the brain is under stress or when coping levels are exceeded.

Dr. Ian Kelleher, RCSI Psychiatry Research Lecturer and study lead, commented: "Our research shows that if we can understand and treat the factors associated with these perceptual abnormalities, we could prevent at least a quarter of suicide attempts and deaths. Given that about 1 million people die by suicide every year, that's a very encouraging prospect for [suicide prevention](#)."

"These findings show the need both for clinicians to pay particular attention to patients reporting experience of psychotic experience, and

for greater funding for research into recognising a psychosis subtype of suicide", added Dr. Kelleher.

Kathryn Yates, RCSI Psychiatry and study co-author, said: "If we are to understand suicide, we need to understand a lot more about perceptual abnormalities—what causes people to hear voices? How do these experiences relate to the biological and social factors involved in suicide risk? There are still a lot of unanswered questions; but this research points to new avenues to improve prediction of suicidal behaviour."

The paper, Psychotic experiences and risk of suicidal ideation, [suicide attempts](#) and suicide deaths: a [systematic review](#) and meta-analysis of longitudinal population studies, is published in *JAMA Psychiatry*.

**More information:** Kathryn Yates et al, Association of Psychotic Experiences With Subsequent Risk of Suicidal Ideation, Suicide Attempts, and Suicide Deaths, *JAMA Psychiatry* (2018). [DOI: 10.1001/jamapsychiatry.2018.3514](#)

Provided by RCSI

Citation: Research reveals that 1 in 4 suicide attempts are associated with perceptual difficulties (2018, November 28) retrieved 6 May 2024 from <https://medicalxpress.com/news/2018-11-reveals-suicide-perceptual-difficulties.html>

|  |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|