

Stress-induced effects on heart blood flow differ for men versus women

November 26 2018

Some patients with coronary artery disease have inadequate blood flow to the heart muscle during periods of mental/emotional stress. This condition—called "mental stress-induced myocardial ischemia" (MSIMI) - is related to the severity of plaque buildup in the coronary arteries in men but not women, reports a study in *Psychosomatic Medicine: Journal of Biobehavioral Medicine*, the official journal of the American Psychosomatic Society.

Mental stress-induced [myocardial ischemia](#) occurs in both men and women, although [younger women](#) tend to have more of this phenomenon. While MSIMI is linked to worse cardiovascular outcomes in both [sexes](#), there may be sex differences in the mechanism through which stress can trigger MSIMI, according to new research by Viola Vaccarino, Md, Ph.D., and colleagues, of Emory University School of Public Health and School of Medicine, Atlanta.

Heart Response to Stress Linked to Coronary Plaque Severity in Men, but Not Women

The researchers examined the sex-specific association between coronary plaque buildup (obstructive coronary disease) assessed from angiograms and MSIMI in 276 participants—141 men and 135 women—with recent myocardial infarction (heart attack). The study targeted young and middle-aged adults (under age 61). All participants underwent conventional stress testing to evaluate myocardial blood flow in response

to a physical or pharmacological challenge (either an exercise stress test or a pharmacological stress test) .

The patients also underwent a [mental stress](#) test, which evaluated myocardial blood flow in response to a stressful situation (a public speaking task). Based on a reduction in myocardial blood flow during this stressful challenge, 17 percent of patients were classified as having MSIMI.

Women had a higher rate of MSIMI (20 percent) than men (15 percent). Twenty-seven percent of patients had conventional stress-induced myocardial ischemia (CSIMI). Only 10 percent of patients had both MSIMI and CSIMI.

In both men and women, the presence of CSIMI was associated with greater obstructive coronary disease severity. For each one-unit increase in severity score, the likelihood of CSIMI increased by about 50 percent, after adjustment for other factors.

In contrast, the presence of MSIMI was associated with more severe plaque buildup only in men: for each one-unit increase in severity score, the likelihood of MSIMI nearly doubled. For women, MSIMI was unrelated to obstructive coronary disease severity.

"There is growing evidence that psychosocial stressors may promote the development and progression of cardiovascular disease in susceptible individuals," according to the authors. However, the relationship between MSIMI and obstructive coronary disease, which means blockages in the major coronary arteries supplying blood to the [heart muscle](#), has been unclear. Women are more likely than men to develop myocardial ischemia without coronary obstruction.

The results support the hypothesis that MSIMI may develop via different

mechanisms in women versus men. Consistent with previous findings on sex differences in heart disease, it may be that MSIMI in women mainly reflect abnormalities other than blockage of the coronary arteries, such as malfunction of the smaller coronary blood vessels (microcirculatory dysfunction).

The sex differences may have important implications for understanding sex-specific vascular effects of psychological stress in general and the mechanisms of MSIMI in particular, since most previous studies of this issue have focused on men. Dr. Vaccarino and colleagues conclude, "These results suggest that MSIMI must be driven by alternative mechanisms especially among [women](#), and provide motivation for further research to understand sex-specific mechanisms for the effects of mental stress on myocardial ischemia and long-term outcomes."

More information: Zakaria Almuwaqqat et al. Sex-Specific Association between Coronary Artery Disease Severity and Myocardial Ischemia Induced by Mental Stress, *Psychosomatic Medicine* (2018).
[DOI: 10.1097/PSY.0000000000000636](https://doi.org/10.1097/PSY.0000000000000636)

Provided by Wolters Kluwer Health

Citation: Stress-induced effects on heart blood flow differ for men versus women (2018, November 26) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-11-stress-induced-effects-heart-blood-differ.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--