

# Weight during adolescence may affect pancreatic cancer risk in adulthood

November 12 2018

---



Axial CT image with i.v. contrast. Macrocystic adenocarcinoma of the pancreatic head. Credit: public domain

New research has linked adolescent obesity with up to a four-fold increased risk of pancreatic cancer later in life. The study's results also suggest that overweight and even higher weight within the "normal"

weight range in men may increase pancreatic cancer risk in a graded manner. The findings are published early online in *Cancer*.

Pancreatic cancer is the sixth most common cause of cancer-related deaths in the world, and studies have linked adult obesity with an increased risk for its occurrence. To uncover any potential associations with adolescent [weight](#), Zohar Levi, MD, of Rabin Medical Center and Tel Aviv University, and his colleagues analyzed 1,087,358 Israeli Jewish men and 707,212 Jewish women who underwent a compulsory physical examination between the ages of 16 and 19 years from 1967 to 2002. Pancreatic cancer incidence through 2012 was identified by linkage to the Israeli National Cancer Registry.

Over a median of 23.3 years of follow up, 551 new cases of [pancreatic](#) cancer cases were identified, including 423 cancers among men and 128 cancers among women. Compared with normal weight (5th to

Citation: Weight during adolescence may affect pancreatic cancer risk in adulthood (2018, November 12) retrieved 27 April 2024 from <https://medicalxpress.com/news/2018-11-weight-adolescence-affect-pancreatic-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------