

Women who give birth to boys much more likely to have postnatal depression

November 6 2018



Credit: CC0 Public Domain

A University of Kent study into postnatal depression (PND) found the odds of developing this condition increased by 79% when mothers had baby boys compared to baby girls.

Overall the researchers identified that women who give birth to males are 71-79% more likely to develop PND. Furthermore, women whose births had complications were 174% more likely to experience PND compared to those women who had no complications.

As a result of their findings, Dr. Sarah Johns and Dr. Sarah Myers in the University's School of Anthropology and Conservation (SAC), conclude that recognising that both male infants and birth complications are PND risk factors should help health professionals in identifying and supporting women who may be more likely to develop this condition.

Their research also showed that while women with a tendency towards symptoms of depression, anxiety, and stress were always at increased risk of PND, they had reduced odds of developing PND after experiencing birth complications. This is likely because these women may receive greater post-birth support because their [mental health concerns](#) were previously recognised. This finding suggests interventions to support women can be effective in preventing PND developing.

Dr. Johns said: 'PND is a condition that is avoidable, and it has been shown that giving women at risk extra help and support can make it less likely to develop. The finding that having a baby boy or a difficult birth increases a woman's risk gives health practitioners two new and easy ways to identify women who would particularly benefit from additional [support](#) in the first few weeks and months'

Dr. Johns and Dr. Myers decided to assess whether there was a relationship between the sex of infants and PND because of the known link between inflammatory immune response and the development of depressive symptoms.

Both the gestation of male foetuses and the experience of [birth](#) complications have documented associations with increased

inflammation, yet, until this study, their relationships with PND were unclear.

Many known risk factors for [depressive symptoms](#) are associated with activation of inflammatory pathways, opening up the potential for identifying new risk factors based on their inflammation causing effects—an idea supported by this study.

The study used complete reproductive histories of 296 [women](#) from contemporary, low fertility populations.

More information: Sarah Myers et al, Male infants and birth complications are associated with increased incidence of postnatal depression, *Social Science & Medicine* (2018). [DOI: 10.1016/j.socscimed.2018.10.008](#)

Provided by University of Kent

Citation: Women who give birth to boys much more likely to have postnatal depression (2018, November 6) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-11-women-birth-boys-postnatal-depression.html>

| |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|