


Why women get PMS and why some are more affected

November 19 2018, by Jayashri Kulkarni



Credit: AI-generated image ([disclaimer](#))

Women have been menstruating throughout history. So it's curious the earliest documented record of what we now know to be premenstrual syndrome (PMS) appeared pretty late in the game. In 1931, psychoanalyst [Karen Horney](#) described increased tension, irritability, depression and anxiety in the week preceding menstruation in one of her

patients.

Now it's [generally accepted](#) up to 80% of [women](#) in their reproductive years experience some PMS. The condition [includes symptoms](#) such as fatigue, poor coordination, feeling out of control, feeling worthless and guilty, headache, anxiety, tension, aches, irritability, [mood swings](#), weight gain, food cravings, no interest in usual activities, cramps, feeling sad or depressed, breast tenderness, sleep problems, and difficulty concentrating.

Premenstrual syndrome is different to premenstrual dysphoric disorder (PMDD), which is rarer (only [3-5% of women](#) of reproductive age experience it) and is listed in the [diagnostic manual](#) of mental disorders. People who experience PMDD have severe depression which is often accompanied by suicidal thoughts. Their onset and offset usually coincide with the premenstrual cycle. Unlike PMS, the severely depressed mood of PMDD usually comes on suddenly.

Reproductive hormones – oestrogen, progesterone and testosterone – are also potent brain hormones. They [influence the brain chemicals](#) responsible for our thoughts, behaviours and emotions. Their amounts fluctuate throughout the menstrual cycle, so the connection between them and mental health is clear. And we are learning more about why some women may be more affected than others.

Brain chemicals and PMS

There is no single clear theory yet to explain exactly which hormones trigger particular chemicals or why only some women experience PMDD or PMS.

But we know some women are susceptible to mood changes due to small fluctuations in reproductive hormones. In these vulnerable women, small

changes in oestrogen and progesterone levels lead to shifts in central brain chemicals (GABA, serotonin and dopamine) that then affects mood and behaviour.

At the same time, many of the physical PMS symptoms such as breast tenderness, bloating, headaches and constipation are a [direct effect](#) of reproductive hormones. So both mind and body are affected.

Oestrogen [appears to be](#) a "protective" hormone, which can [improve psychotic symptoms](#) (such as those common in schizophrenia) as well as depression. Oestrogen directly influences the neurotransmitters serotonin and dopamine to achieve this positive effect.

So depression and other adverse mental symptoms can appear or worsen during phases when oestrogen is low. This happens during the four to seven days before menstruation, and during the transition into menopause.

Progesterones can have the opposite effect. Many women who take a progesterone-only contraceptive pill (the mini-pill) [experience depression](#). There are certain types of progesterone in the combined oral contraceptive pill that can be very depressive.

What about the more severe symptoms?

Recent [work suggests](#) PMDD is the result of brain neurochemicals responding in unusual ways to fluctuations in brain oestrogen, progesterone and testosterone, as well as the hormones released by the pituitary gland that determine the levels and fluctuations of these [reproductive hormones](#).

Other [studies](#) about the cause of PMDD reveal that a breakdown product of progesterone – called allopregnanolone (ALLO) – is a critical

stimulator of a receptor on a part of the GABA transmitter. When stimulated, the GABA system can alleviate anxiety. Benzodiazepine drugs like diazepam (Valium) stimulate the GABA system and help to calm down agitation.

In this way, ALLO works as an "anti-anxiety" hormone. Just like oestrogen, progesterone levels (and its metabolite ALLO levels) fall in the premenstrual phase.

Women who have PMDD are often agitated, anxious and depressed during the premenstrual phase. A [newer theory](#) is that their brain chemistry isn't reacting normally to ALLO, so they become anxious. This is important to explore further and already new drugs that impact ALLO are being developed and tested.

PMDD is complex, like many mental health conditions, and there is an interplay between psychological and social issues as well as hormonal and neurochemical factors. Tertiary education, supportive relationships, fewer socioeconomic struggles and good physical health [appear to be helpful](#), but do not mitigate PMDD completely. Overall, PMDD appears to be biologically driven.

How can we treat it?

Understanding the body-mind connections in both PMS and PMDD is critical for developing effective management strategies for the many women who suffer from significant depression and other issues every month.

Management options need to consider all aspects of the woman's life including her work, relationships, past traumas, current physical health and daily demands. Many women experiencing PMDD require [hormone](#) treatment and other strategies such as antidepressant medication to help

them improve their quality of life.

It's a good idea for women experiencing PMDD or PMS to keep a diary of their cycles and moods. Women can be reassured their observations connecting hormones and moods are valid. It is important women with PMS/PMDD seek help from health professionals who will explore specific targeted treatments with them. Above all, it is important to recognise the links between hormones and [mental health](#).

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Why women get PMS and why some are more affected (2018, November 19) retrieved 5 May 2024 from <https://medicalxpress.com/news/2018-11-women-pms-affected.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--