


Breast cancer drugs could help treat resistant lung cancers

December 26 2018



Lung CA seen on CXR. Credit: [CC BY-SA 4.0](https://creativecommons.org/licenses/by-sa/4.0/) James Heilman, MD/Wikipedia

A class of drugs used to treat certain breast cancers could help to tackle lung cancers that have become resistant to targeted therapies, suggests a new study in mice from the Francis Crick Institute and the Institute of Cancer Research (ICR).

The research, published in *Cell Reports*, found that [lung](#) tumours in mice caused by mutations in a gene called EGFR shrunk significantly when a protein called p110 α was blocked.

Drugs to block p110 α are currently showing promise in [clinical trials](#) against certain breast cancers, so could be approved for clinical use in the near future. The new findings suggest that these drugs could potentially benefit patients with EGFR-mutant lung cancers whose tumours have become resistant to [treatment](#).

"At the moment, patients with EGFR-mutant lung cancers are given targeted treatments that are very effective for the first few years," explains study leader Professor Julian Downward, who has labs at the Crick and the ICR. "These drugs are improving, but unfortunately after a couple of years the [cancer](#) usually becomes resistant and starts to grow and spread again. The second line of treatment is currently conventional chemotherapy, which is not targeted and has substantial side-effects.

"Our new study suggests that it would be worth investigating whether p110 α inhibitors could be used as a second-line therapy. As our research is at such an [early stage](#), more research in mice and patient cells would be needed before even considering clinical trials, but it opens up a promising avenue of investigation."

For this research, the team targeted a specific interaction between the RAS protein and p110 α . The RAS gene is mutated in around one in five cancers, causing uncontrolled growth, and is a key focus of Julian's research. When they blocked this interaction in genetically modified

mice with EGFR mutations, their tumours shrank significantly.

Before the intervention, the tumours filled around two thirds of the space inside the lung. When the interaction between RAS and p110 α was genetically blocked, this shrank significantly to about a tenth of the space inside the lung. The intervention also had very few side-effects.

"As we wanted to pinpoint the specific interaction responsible, we used a genetic technique that would not be practical in a patient treatment," says Julian. "We're looking to develop ways to do this with drugs, as blocking this specific pathway would significantly reduce side-effects, but this work is many years from the clinic. In the medium-term, investigating existing drugs that inhibit p110 α will be the next step. While these have side-effects, including temporary diabetes-like symptoms during treatment, they are still less toxic than chemotherapy."

Provided by The Francis Crick Institute

Citation: Breast cancer drugs could help treat resistant lung cancers (2018, December 26)
retrieved 28 April 2024 from
<https://medicalxpress.com/news/2018-12-breast-cancer-drugs-resistant-lung.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--