

How does diet during pregnancy impact allergies in offspring?

December 14 2018



Karen Robbins, M.D., an allergist at Children's National Health System. Credit: Children's National Health System

A small percentage of women said they consumed fewer allergens during pregnancy to stave off food allergies in their newborns, according to preliminary research Karen Robbins, M.D., presented during the American College of Asthma Allergy and Immunology 2018 Annual

Scientific Meeting.

Pregnant [women](#) routinely swear off alcohol and tobacco to boost their chances of having a healthy baby. What about common [food](#) allergens like nuts and milk?

There are scant data that describe how often pregnant women deliberately stop eating a specific food item in order to prevent future food allergies in their newborns. As a first step toward addressing this data gap, a research team led by Karen Robbins, M.D., an allergist at Children's National Health System, pored through a [longitudinal study](#) conducted by the Food and Drug Administration (FDA) and the Centers for Disease Control and Prevention.

About 4,900 pregnant women completed the Infant Feeding Practices Study II prenatal questionnaire from May 2005 to June 2007. The study tracked 2,000 pregnant women from the third trimester of pregnancy and their infants through the first year of life. A small percentage of women said they had consumed fewer allergens during pregnancy to stave off food allergies in their newborns, according to a poster Dr. Robbins presented during the American College of Asthma Allergy and Immunology 2018 Annual Scientific Meeting. While their numbers were small, most of these women reported giving up major allergens like nuts, milk or eggs during pregnancy, including:

- 144 (2.9 percent) reported restricting their diet in some way to prevent future food allergies in their offspring
- 84 women (1.7 percent) ate fewer nuts
- 15 women (.3 percent) ate fewer eggs and
- 2 women (.04 percent) ate/drank consumed less dairy/milk.

"At the time the survey was conducted, few pregnant women in this large data set said they gave up certain foods with the express aim of avoiding

a food allergy in their babies," Dr. Robbins says. "However, mothers who had an older child with a food allergy or who had food allergies themselves had significantly higher odds of trying this food avoidance strategy."

Despite the diet changes, infants born to these [expectant mothers](#) were twice as likely to experience problems with food at age 4 months—though not at age 9 months or 12 months. And these infants were no more likely to be diagnosed with a food allergy.

According to the FDA, millions of Americans suffer a food allergy each year. Reactions can range from mild to life-threatening and can begin soon after eating a problematic food item or an ingredient from that food. Among the most common allergenic foods are milk, eggs, fish, shellfish, tree nuts, peanuts, wheat and soybeans.

"We really need to know more about how often targeted food avoidance occurs among U.S. [pregnant women](#) who have a family history of food allergies," Dr. Robbins adds. "We hope to learn what factors into these women's decision-making as well as why many of them settled on food avoidance as a potential strategy to try to prevent food [allergy](#) in their infants."

More information: "Prenatal food allergen avoidance practices for food allergy prevention" American College of Asthma Allergy and Immunology 2018 Annual Scientific Meeting

Provided by Children's National Medical Center

Citation: How does diet during pregnancy impact allergies in offspring? (2018, December 14) retrieved 23 April 2024 from

<https://medicalxpress.com/news/2018-12-diet-pregnancy-impact-allergies-offspring.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.