

Kicking, yelling during sleep? Study finds risk factors for violent sleep disorder

December 26 2018



Credit: CC0 Public Domain

Taking antidepressants for depression, having post-traumatic stress disorder or anxiety diagnosed by a doctor are risk factors for a disruptive and sometimes violent sleep disorder called rapid eye movement (REM)

sleep behavior disorder, according to a study published in the December 26, 2018, online issue of *Neurology*, the medical journal of the American Academy of Neurology. The study also found men are more likely to have the disorder.

REM sleep is the dream state of sleep. During normal REM sleep, your brain sends signals to prevent your muscles from moving. However, for people with REM sleep behavior disorder, those signals are disrupted. A person may act out violent or action-filled dreams by yelling, flailing their arms, punching or kicking, to the point of harming themselves or a sleep partner.

"While much is still unknown about REM sleep behavior disorder, it can be caused by medications or it may be an early sign of another neurologic condition like Parkinson's disease, dementia with Lewy bodies or multiple system atrophy," said study author Ronald Postuma, MD, MSc, of McGill University in Montreal, Canada, and a member of the American Academy of Neurology. "Identifying lifestyle and personal risk factors linked to this sleep disorder may lead to finding ways to reduce the chances of developing it."

The study looked at 30,097 people with an average age of 63. Researchers screened participants for a variety of health conditions and asked about lifestyle, behavior, social, economic and psychological factors.

In addition, every participant was asked, "Have you ever been told, or suspected yourself, that you seem to act out your dreams while asleep?"

Researchers then identified 958 people, or 3.2 percent, with possible REM sleep behavior disorder, after excluding participants with Parkinson's disease, dementia, Alzheimer's disease or sleep apnea.

Researchers found those with the disorder were over two-and-a-half times as likely to report taking antidepressants to treat depression, with 13 percent of those with the disorder taking them compared to 6 percent of those without the disorder. People with the disorder were also two-and-a-half times as likely to have post-[traumatic stress disorder](#). They were twice as likely to have [mental illness](#), and over one-and-a-half times as likely to have psychological distress.

Other findings were that men were twice as likely as women to have possible REM sleep behavior disorder; 59 percent of those with the disorder were male, compared to 42 percent of those without the disorder. People with possible REM sleep behavior disorder were 25 percent more likely than those without the disorder to be moderate to heavy drinkers, with 19 percent of those with the disorder moderate to heavy drinkers compared to 14 percent of those without the disorder. They had slightly less education, an average of 13.2 years of education compared to an average of 13.6 years for those without the disorder. They also had lower income and were more likely to have smoked.

"Our research does not show that these [risk factors](#) cause REM sleep behavior disorder, it only shows they are linked," said Postuma. "Our hope is that our findings will help guide [future research](#), especially because REM sleep behavior disorder is such a strong sign of future neurodegenerative disease. The more we understand about REM sleep behavior disorder, the better positioned we will be to eventually prevent neurologic conditions like Parkinson's disease."

A limitation of the study was that 96 percent of participants were white, meaning the results may not apply to people of other ethnic backgrounds.

More information: [Abstract/Full Text \(subscription or payment may be required\)](#)

Provided by American Academy of Neurology

Citation: Kicking, yelling during sleep? Study finds risk factors for violent sleep disorder (2018, December 26) retrieved 18 April 2024 from <https://medicalxpress.com/news/2018-12-factors-violent-disorder.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.