

Control HIV by treating schistosomiasis, new study suggests

December 13 2018



Schistosome Parasite: This schistosome parasite enters the body through the skin of persons coming in contact with infested waters. The adult worm lives in the veins of its host. The parasite is magnified x256 in this photograph. Credit:

Wetzel and Shaefer, Wikimedia Commons, Public Domain, 2001

Of the 34 million people worldwide with HIV, and the 200 million with schistosomiasis, the majority live in Africa— where millions of people are simultaneously infected with both diseases. Now, researchers reporting in *PLOS Neglected Tropical Diseases* have shown that schistosomiasis infections are associated with increased HIV onward transmission, HIV acquisition in HIV negative women with urogenital schistosomiasis, and progression to death in HIV positive women.

Schistosomiasis is caused by a [parasitic worm](#) and is second only to malaria in terms of parasitic diseases with the most global health impact. Schistosomiasis may cause lesions in the [genital tract](#) and has been shown—in cross-sectional studies— to be associated with HIV prevalence.

In the new work, Dr. Kristin Wall, of Emory University, USA, and colleagues retrospectively tested the blood sera of 2,145 people enrolled in a cohort of HIV-discordant heterosexual couples in Zambia. Each sample was measured for schistosome-specific antibody levels. In addition, health information including HIV status over time and death was available for the cohort.

59% of all participants were positive for a schistosome-specific antibody response. Among men and women positive for HIV, baseline response for schistosomiasis was associated with an increased risk of transmitting HIV to their uninfected partner. In addition, among women, schistosomiasis was associated with increased acquisition of HIV (in HIV negative women with urogenital schistosomiasis) and increased progression to death (in HIV positive women).

"Since treatment of schistosome infections with praziquantel is inexpensive, effective, and safe, [schistosomiasis](#) prevention and treatment strategies may be a cost-effective way to reduce not only the symptoms associated with the infection, but also new cases of HIV and death among HIV+ persons," the researchers say.

More information: Wall KM, Kilembe W, Vwalika B, Dinh C, Livingston P, et al. (2018) Schistosomiasis is associated with incident HIV transmission and death in Zambia. *PLOS Neglected Tropical Diseases* 12(12): e0006902. doi.org/10.1371/journal.pntd.0006902

Provided by Public Library of Science

Citation: Control HIV by treating schistosomiasis, new study suggests (2018, December 13) retrieved 26 April 2024 from <https://medicalxpress.com/news/2018-12-hiv-schistosomiasis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--