

Insufficient nutrition during fetal development may lead to early menopause

December 5 2018

Previous studies have demonstrated that fetal malnutrition can lead to adult chronic disorders such as type 2 diabetes and coronary artery disease. A new study out of China now suggests that it also can lead to early menopause and premature ovarian failure. Results are published online today in *Menopause*, the journal of The North American Menopause Society (NAMS).

Infants are especially sensitive to changes in their environment while still in the womb, during their earliest stages of development. It has already been documented that the development of the hypothalamic-pituitary-gonadal axis during the fetal stage plays a critical role in adulthood [reproductive health](#). Natural menopause is a milestone of ovarian aging that results in the end of a woman's reproductive years.

Although several studies have investigated the association between famine exposure in [early life](#) and risk of various metabolic diseases in adulthood, the association with reproductive aging was not evaluated. This new study involving nearly 2,900 Chinese women specifically sought to address the effect of early life exposure to famine on age at menopause.

The study concluded that prenatal famine was associated with a higher risk of early menopause (age younger than 45 years), as well as a higher risk of premature ovarian failure. Although [study participants](#) were born during China's infamous famine occurring between 1956 and 1964, the study provides valuable insights into the benefits of proper nutrition

during early life stages for women of any culture.

Study results appear in the article "Early life exposure to famine and reproductive aging among Chinese women."

"The findings that natural menopause occurs earlier after prenatal famine exposure suggests that food deprivation during early fetal life affects how long the future ovaries function," says Dr. JoAnn Pinkerton, NAMS executive director. "For those women, if they are not taking estrogen therapy until the average age of menopause, their [early menopause](#) could be associated with increased risk of heart disease, osteoporosis, depression, and memory changes and changes in vaginal and sexual health."

Provided by The North American Menopause Society

Citation: Insufficient nutrition during fetal development may lead to early menopause (2018, December 5) retrieved 25 April 2024 from <https://medicalxpress.com/news/2018-12-insufficient-nutrition-fetal-early-menopause.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--