

# Key signs and symptoms to recognise when terminally ill patients are close to death

December 11 2018

---



Credit: CC0 Public Domain

Research jointly undertaken by Kingston University and St George's, University of London's new Associate Dean of Research and Enterprise has identified the key signs and symptoms expert doctors use to

recognise when terminally ill patients are close to death.

Professor Priscilla Harries, who is also the Director of the Centre of Health and Social Care Research in the Faculty of Health, Social Care and Education, run jointly by Kingston and St George's, co-authored the study led by researchers at the Marie Curie Palliative Care Research Department at University College London.

As part of the research a group of palliative [care](#) doctors, who had particular expertise in recognising when a patient is nearing death, were asked to predict whether a patient was close to death in 20 real-life cases. Those with the most accurate results were given an additional 50 hypothetical cases and asked to estimate the probability of death within the next 72 hours.

Through this exercise, Professor Harries and her co-researchers were able to determine the relative importance of different factors behind the doctors' survival predictions. The most influential was the Palliative Performance Score, which measures how much support someone needs based on their functional ability. The second most important factor was the presence of an abnormal pattern of breathing called Cheyne-Stokes respiration.

The research, published in the Journal *BMJ Open*, followed a previous study that found some doctors were frequently inaccurate when predicting how long those living with terminal illnesses will survive.

It is hoped the study, titled "How do Palliative Care Doctors Recognise Imminently Dying Patients? A Judgement Analysis," will help inform the development of educational training materials to teach novice doctors, medical students and other [healthcare professionals](#) how to model their own judgement policies on those of the experts.

The researchers are now developing and testing online education through a randomised controlled trial. "If effective, it will enable us to teach novice [doctors](#) and [medical students](#) how to model their own judgement policies on those of the experts," Professor Harries said.

"It is so important to have this last stage of life recognised—for patients to be treated appropriately, to move home if wished, and to notify family and loved ones so they have the chance to say goodbye. Dying well is an important stage of life and should be a key aspect of the high quality health services we provide," she added.

**More information:** Nicola White et al. How do palliative care doctors recognise imminently dying patients? A judgement analysis, *BMJ Open* (2018). [DOI: 10.1136/bmjopen-2018-024996](https://doi.org/10.1136/bmjopen-2018-024996)

Provided by St. George's University of London

Citation: Key signs and symptoms to recognise when terminally ill patients are close to death (2018, December 11) retrieved 24 April 2024 from <https://medicalxpress.com/news/2018-12-key-symptoms-recognise-terminally-ill.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--