

Q&A: Deep brain stimulation may reduce seizures caused by epilepsy

December 5 2018, by From Mayo Clinic News Network



Credit: CC0 Public Domain

Dear Mayo Clinic: How does deep brain stimulation for epilepsy work? Who's a good candidate for this treatment? Is it effective?



A: Deep <u>brain</u> stimulation is a technique that uses a wire placed permanently in the brain to send electrical pulses to the brain. It has been approved by the Food and Drug Administration to treat <u>epilepsy</u> that hasn't responded to other forms of therapy. In most people, <u>deep brain stimulation</u> doesn't completely eliminate seizures caused by epilepsy, but it can significantly reduce seizures.

Epilepsy is a central nervous system disorder. In a person with epilepsy, nerve cell activity in the brain becomes abnormal, causing seizures and sometimes loss of consciousness. The symptoms of a seizure can vary widely from one person to another. For example, some people with epilepsy stare blankly for a few seconds during a seizure or appear confused. Others may lose consciousness and have repetitive jerking of their arms and legs.

Even mild epilepsy requires treatment because seizures can be dangerous during activities such as driving. Medication to reduce or eliminate seizures usually is the first step in treatment. For about two-thirds of people with epilepsy, seizures are effectively controlled with the first or second anti-seizure drug they try.

When medication doesn't provide adequate seizure control, surgery may be an option. Epilepsy surgery typically involves removing the area of the brain that's causing seizures. That approach only works, however, when the place within the brain that's triggering the seizures can be identified clearly. In some people, that's not possible. It's those individuals who are most likely to benefit from deep brain stimulation.

Deep brain stimulation for epilepsy involves one surgery where you are asleep. During surgery, the surgeon implants a thin wire lead with several contacts, or electrodes, at the tip into an area deep within the brain called the "thalamus." The thalamus is basically a relay station that distributes signals that come from the body's senses to other regions of the brain.



After an imaging exam confirms that the electrodes are placed properly, a battery-operated device called a pulse generator is implanted under the skin near the collarbone. Wires running from the electrodes in the brain are placed under the skin and connected to the generator. The generator is programmed to send continuous electrical pulses to the brain.

A large clinical research trial found that in people with epilepsy whose seizures didn't respond to other therapies, around 15 percent became seizure-free for more than six months after deep brain stimulation. Although that number is fairly low, it represents significant—often life-changing—improvement for those individuals.

But the goal of deep brain stimulation typically is not complete relief of all seizures. Instead, it's used as a method to reduce the number of seizures a person has. The same clinical trial showed that between 50 and 60 percent of patients in the study had a decrease in their seizures in response to deep brain stimulation. The research also showed that within the group who did respond, the number of seizures they had continued to decrease over time.

Although deep brain stimulation is a major surgical procedure, the risks involved are fairly low, compared to other types of brain surgery. And if deep brain stimulation is found to be ineffective for a patient, it's reversible. The electrodes and generator can be removed without injuring the brain. That makes it an attractive option to consider for people with epilepsy who haven't had success with other treatment and in whom the location within the brain where the <u>seizures</u> arise cannot be pinpointed.

©2018 Mayo Foundation for Medical Education and Research Distributed by Tribune Content Agency, LLC.

Citation: Q&A: Deep brain stimulation may reduce seizures caused by epilepsy (2018, December



5) retrieved 23 April 2024 from https://medicalxpress.com/news/2018-12-qa-deep-brain-seizures-epilepsy.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.