

Stress in new mothers causes lasting health risks, depending on race, ethnicity, poverty

December 14 2018

African-American women undergo more physical "wear-and-tear" during the first year after giving birth than Latina and white women, a consequence that may have long-lasting health effects, according to a study of a diverse group of more than 2,400 low-income women.

The study in today's (Friday, Dec. 14) *American Journal of Perinatology* involved [women](#) of diverse racial and ethnic backgrounds who were interviewed and evaluated at five different clinical sites in the United States.

In addition to insight into [health risks](#) facing [new mothers](#), researchers united by the Eunice Kennedy Shriver National Institute of Child Health and Human Development's Community Child Health Network found evidence that women who breastfeed their babies may receive some protection from the health damage caused by physical stress of pregnancy and giving birth.

"All mothers are affected by stress, but [low-income women](#) and especially African-American and Hispanic women have more adverse health-risk profiles during their children's first years of life," said Sharon Landesman Ramey, a professor and distinguished research scholar at the Fralin Biomedical Research Institute at VTC and an author of the paper. "Our study was designed to look for biomarkers that are sensitive to psychological and physical stressors, and in turn determine whether those stressors contribute to poor outcomes for mothers and children."

Working groups of clinicians guided by scientists and [community members](#) recorded blood pressures, heart rates, cholesterol profiles, body mass indexes, waist-hip ratios, and other biomarkers in women six months and one year after they had given birth. In addition, women were assessed before they gave birth, some even before they became pregnant.

The readings were used to create a composite measurement of "allostatic load," which represents the cumulative physical and psychological strain on their bodies after delivery. The study may be the first to examine a large group of women at intervals over the course of a year using demographic, biometric, and biomarker measures.

Readings were taken to look at the post-delivery effects of pregnancy on mothers' cardiometabolic risks in partnership with groups at five clinical sites in the U.S., including medical centers in Chicago, Baltimore, Los Angeles, Washington, D.C., and North Carolina.

More than one half of the women in the study were classified as overweight or obese prior to becoming pregnant.

"We thoroughly looked at the effects of stress alongside levels of poverty and [minority status](#) to understand poor health outcomes for moms and children," said Madeleine Shalowitz, a Research Professor of Pediatrics at the Pritzker School of Medicine at the University of Chicago, Program Director of Outcomes Research at NorthShore University HealthSystem Research Institute, and a co-investigator in the study. "Filling this knowledge gap could lead to health interventions to lower the risk of chronic disease for mothers, many of whom are planning to have more children. As many of these women will become pregnant again, improving their health will lead to a healthier next pregnancy."

During pregnancy, dramatic changes occur in a woman's immune and

cardiovascular systems to support the developing fetus.

In healthy women, maternal physiology gradually returns to normal within a year after delivery, but a persistently elevated allostatic load increases a woman's risk for chronic diseases across a lifetime.

The researchers said the results of the study are important for improving health and also to address interventions for disadvantaged groups.

As for why breastfeeding was shown to provide some [health benefits](#) 12 months postpartum, the study said it could be a cumulative effect reflecting longer durations of breastfeeding compounded by a better socioeconomic status for women who can afford to breastfeed for longer periods.

A large research network in the National Institutes of Health united with community members to execute the research.

"This study was extraordinary because it included community members as full partners at all of the clinical sites, working alongside health-care providers and scientists to determine how stress is affecting mothers and the next generation of babies," Landesman Ramey said. "The hope is that when people realize health-care providers, citizen activists, and neighbors approved the work and contributed to it, they will have more confidence in the information and it will be shared in a good way, to improve [health](#) outcomes."

Provided by Virginia Tech

Citation: Stress in new mothers causes lasting health risks, depending on race, ethnicity, poverty (2018, December 14) retrieved 10 April 2024 from <https://medicalxpress.com/news/2018-12-stress-mothers-health-ethnicity-poverty.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.