

Continuity of care tied to lower expenditures, hospitalization

December 12 2018



(HealthDay)—Continuity of care scores are significantly associated with



lower expenditures and hospitalization rates, according to a study published in the November/December issue of the *Annals of Family Medicine*.

Andrew Bazemore, M.D., M.P.H., from the Robert Graham Center for Policy Studies in Washington, D.C., and colleagues used Medicare claims data for 1,448,952 beneficiaries obtaining care from a nationally representative sample of 6,551 primary care physicians to calculate four physician-level, claims-based continuity measures. To create physician-level scores, researchers averaged patient-level continuity scores attributed to a single physician. They assessed the correlations between total Medicare Part A and B expenditures and hospitalization.

The researchers found that continuity scores were highly correlated (correlation coefficients, 0.86 to 0.99); for each, greater continuity was correlated with similar outcomes. The adjusted expenditures were 14.1 percent lower for beneficiaries cared for by physicians in the highest versus the lowest quintile of the Bice-Boxerman continuity score (\$6,958 versus \$8,092); between the highest and lowest continuity quintiles, the odds of hospitalization were 16.1 percent lower (odds ratio, 0.839).

"In summary, this study contributes to the overwhelming evidence of the value of continuity care," the authors write. "Continuity is one of a handful of core tenets of primary care that should be incorporated into official primary care measures as we shift from paying for services to paying for value."

More information: Abstract/Full Text

Editorial

Copyright © 2018 HealthDay. All rights reserved.



Citation: Continuity of care tied to lower expenditures, hospitalization (2018, December 12) retrieved 27 April 2024 from

https://medicalxpress.com/news/2018-12-tied-expenditures-hospitalization.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.