


Antidepressant use in seniors linked to risk for hip fracture

January 3 2019



(HealthDay)—Among older adults, antidepressant users sustain more hip



fractures than nonusers both before and after therapy initiation, according to a study published online Jan. 2 in *JAMA Psychiatry*.

Jon Brännström, M.D., from Umeå University in Sweden, and colleagues examined the correlation between antidepressant drug treatment and <a href="https://hip.nc.nih.google.com/hip.google.

The researchers found that in the year before and the year after initiation of therapy, antidepressant users sustained more than twice as many hip fractures as nonusers (2.8 versus 1.1 percent and 3.5 versus 1.3 percent, respectively). Adjusted analyses revealed the highest odds ratios for the correlation between antidepressant use and hip fracture 16 to 30 days before the prescription was filled (odds ratio, 5.76). The highest odds ratios were also seen 16 to 30 days before treatment in subgroup analyses of age groups, men and women, and individual antidepressants, with no clear dose-response relationship.

"For most <u>older adults</u>, the toll of untreated depression will likely outweigh the <u>potential risks</u> associated with antidepressant use, including falls or fractures," write the authors of an accompanying editorial.

More information: Abstract/Full Text (subscription or payment may be required)

Editorial (subscription or payment may be required)

Copyright © 2018 HealthDay. All rights reserved.

Citation: Antidepressant use in seniors linked to risk for hip fracture (2019, January 3) retrieved



3 May 2024 from

https://medicalxpress.com/news/2019-01-antidepressant-seniors-linked-hip-fracture.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.