

How bad will my postpartum depression be in 12 months?

January 16 2019, by Kristin Samuelson



Credit: George Hodan/Public Domain

A new Northwestern Medicine study was able to successfully predict if a new mother would experience worsening depressive symptoms over the first year after giving birth by identifying four maternal characteristics that put her at risk.

Identifying these factors early in the [postpartum period](#) will allow

mothers to seek treatment earlier and improve their chance of a full recovery, the authors said.

The four characteristics included the number of children the mother has; her ability to function in general life, at work and in relationships; her [education level](#), which can determine access to resources; and her [depression severity](#) at four to eight weeks postpartum. The predictions from the study were 72.8 percent accurate.

The study was published January 15 in the journal *Depression & Anxiety*.

"By the time a mother comes in for her six-week postpartum visit, we have the potential to predict the severity of her [depression](#) over the next 12 months," said first author Sheehan Fisher, assistant professor of psychiatry and behavioral sciences at Northwestern University Feinberg School of Medicine. "This would be a game-changer for mothers and their clinicians because we could encourage early intervention so moms have better odds of success with their treatment over time."

A mother with postpartum depression can fall into one of three depression trajectories, ranging from gradual remission (over time she starts to get better), to partial improvement (by 12 months postpartum, she is headed in a positive direction but continues to have symptoms), to chronic severe (her symptoms start at the same level as the partial improvement trajectory but worsen over time).

"It's not just a question of 'Is the mother feeling depressed?' but rather, 'Which way is she headed in her depression?'" Fisher said. "If her depression symptoms are going to get worse over time, she needs to be proactive about treatment."

Fisher hopes the findings will lead to improved step care for [mothers](#) in all three depression trajectories, meaning the level of care can be

tailored to each woman.

Mothers with [postpartum depression](#) typically experience difficulty sleeping, feelings of worthlessness or excessive guilt, difficulty coping with [negative emotions](#), have an inability to focus or concentrate on things and generally feel a lot of emotional distress, Fisher said.

Postpartum depression impacts not only the mother but also can negatively impact her child's functioning and health. It can affect the child's emotional development, ability to regulate their own emotions and confer a higher risk for anxiety and depression.

The longer a woman's depression goes untreated, the more difficult it is to get her back on track, Fisher said. It can also take a while to find the right medication and get access to the right provider.

"It only complicates things if the mother doesn't start her treatment until later on," Fisher said.

Treatment for women in the chronic severe group would differ based on the individual but could include psychotherapy and/or medication, Fisher said. Clinicians might enlist the help of the father or other family members or might seek high-level care like an intensive outpatient course of treatment for the mother.

The [longitudinal study](#) looked at data collected between 2006 and 2011 of women delivering at an academic medical center in Pittsburgh, Pa. Women with a postpartum [depressive disorder](#) participated and completed symptom severity assessments at four-to-eight weeks (intake), three months, six months and 12 months postpartum. Clinicians interviewed the women about the severity of their [depressive symptoms](#), medical and psychiatric history, functioning, obstetric experience and infant status.

The scientists determined a woman's score based on the four characteristics and, using a computational algorithm that predicts her depression trajectory, provided the odds of which group the woman would fall in.

More information: Sheehan D. Fisher et al, Four maternal characteristics determine the 12-month course of chronic severe postpartum depressive symptoms, *Depression and Anxiety* (2019). [DOI: 10.1002/da.22879](https://doi.org/10.1002/da.22879)

Provided by Northwestern University

Citation: How bad will my postpartum depression be in 12 months? (2019, January 16) retrieved 20 March 2024 from <https://medicalxpress.com/news/2019-01-bad-postpartum-depression-months.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--