

# How does the brain learn by talking to itself?

January 2 2019

---



Credit: CC0 Public Domain

Humans, like other animals, possess an enormous learning capacity that allows for the apprehension of new sensory information to master new skills or to adapt to an ever-changing environment. However, many of the mechanisms that enable people to learn remain poorly understood. One of the greatest challenges of systems neuroscience is to explain how

synaptic connections change to support adaptive behaviours. Neuroscientists at the University of Geneva (UNIGE), Switzerland, previously showed that synaptic learning mechanisms in the brain's cortex are dependent on feedback from deeper brain regions. They have now deciphered how this feedback gates synaptic strengthening by switching particular inhibitory neurons on and off. This study, published in *Neuron*, not only constitutes an important milestone in the understanding of the mechanisms for perceptual learning, but may also offer insight into computerized learning systems and artificial intelligence.

The cortex – the brain's outer and largest region – is important for higher cognitive functions, complex behaviours, perception and learning. Upon the arrival of a sensory stimulus, the cortex processes and filters its information before it passes the most relevant aspects on to other brain regions. Some of these brain regions, in turn, send information back to the cortex. These loops, known as "feedback systems," are thought to be essential for the functioning of cortical networks and their adaptation to new sensory information. "For [perceptual learning](#) – which is the improved ability to respond to a sensory stimulus – neuronal circuits need to first assess the importance of the incoming sensory information and then refine the way it is processed in the future. Feedback systems to a degree confirm that those synapses that were responsible for transmitting the information to other brain areas did this correctly," explains Anthony Holtmaat, professor in basic neurosciences at the UNIGE Faculty of Medicine, who directed this study.

## **How the whiskers highlight the feedback systems**

The whiskers on a mouse's snout are specialized in tactile sensing and play a major role in the animal's ability to comprehend aspects of its direct environment. The part of the cortex that processes sensory information from the whiskers continuously optimizes its synapses in

order to learn new aspects about the tactile environment. Therefore, it constitutes an interesting model for understanding the role of feedback systems in synaptic learning mechanisms.

The UNIGE scientists isolated a whiskers-related feedback circuit, and used electrodes to measure the electrical activity of neurons in the cortex. They then mimicked the [sensory input](#) by stimulating a specific part of the cortex known for processing this information, and, at the same time, used light to control the feedback circuit. "This ex vivo model allowed us to control the feedback independently from the sensory input, which is impossible to do in vivo. However, disconnecting the sensory input from the feedback was essential to understanding how the interplay between the two leads to synaptic strengthening," adds Holtmaat.

## **Inhibiting neurons gate the information**

The team found that both components, when triggered separately, activate a wide range of neurons. However, when activated simultaneously, some neurons actually decrease their activity.

"Interestingly, the neurons that are inhibited when the sensory input and the feedback occur together usually inhibit neurons that are important for perception, this is known as an inhibition of inhibition or a disinhibition," explains Leena Williams from the UNIGE Faculty of Medicine, the study's first author. "Thus, these neurons act like a gate for the incoming information, and which is normally closed. But when feedback comes in, the gate is opened, allowing those synapses that take care of the primary sensory information to increase their strength. With this study we have identified how feedback possibly optimizes synaptic connections to better prepare for future incoming information," she adds.

Now that they have identified which neurons are involved in this

mechanism, these scientists will test their results in "real life" to check whether the inhibiting [neurons](#) will behave as predicted when a mouse needs to learn new sensory information or when it discovers new aspects in its tactile environment.

## Deep learning: mimicking natural intelligence

How do brain circuits optimize themselves? How can a system teach itself by reading out its own activity? Apart from being relevant to learning in animals, this question is also at the heart of machine learning programs. Indeed, some deep learning specialists try to mimic brain circuits to build artificially intelligent systems. Insights such as provided by the UNIGE team might be relevant for unsupervised learning, a branch of machine learning that occupies itself with circuit models that are able to self-organize and optimize the processing of new information. This is important for the creation of efficient voice or face recognition programmes, for example.

**More information:** Leena E. Williams et al. Higher-Order Thalamocortical Inputs Gate Synaptic Long-Term Potentiation via Disinhibition, *Neuron* (2018). [DOI: 10.1016/j.neuron.2018.10.049](https://doi.org/10.1016/j.neuron.2018.10.049)

Provided by University of Geneva

Citation: How does the brain learn by talking to itself? (2019, January 2) retrieved 23 April 2024 from <https://medicalxpress.com/news/2019-01-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--