

# Cannabis use disorder: The policy climate matters

January 23 2019

---

Adolescents and young adults living in states with more liberal policies reported higher average rates of past-year cannabis use than those in states with more conservative policies, according to a new study conducted at Columbia University Mailman School of Public Health. However, the rates of cannabis use disorder—abuse or dependence on the drug—were significantly lower in states with more liberal policies compared to states with more conservative policies, for ages 12 to 17, and marginally lower for ages 26 and older. These results remained significant even when controlling for the presence of medical cannabis laws. This study is one of the first to assess the relationship between policy liberalism and health outcomes, and specifically cannabis use-related outcomes. The findings are published in the *International Journal of Drug Policy*.

"The majority of existing work has explored the relationship between medical cannabis laws and cannabis outcomes, whereas our results identified important relationships between the state-level [policy](#) context as a whole, and cannabis use outcomes," said Morgan Philbin, Ph.D., assistant professor of Sociomedical Sciences, and first author. "While this research does not suggest that being in a liberal state causes people to use cannabis, or have lower rates of cannabis use disorder, it does highlight how states may differ beyond substance use policies, and how these differences also merit attention."

Using nationally-representative state-level data, the researchers examined the associations between policy liberalism and cannabis use

and cannabis use disorder among past year users. Data were obtained for ages 12-17, 18-25, and 26 and older from the 2004-2006 and 2010-2012 National Survey on Drug Use and Health.

Past year cannabis use was consistently higher in liberal compared to conservative states, and remained significantly higher for ages 12-17 and 18-25 after adjusting for medical cannabis law status. As of December 2018, a total of 33 states had approved medical cannabis laws and eight states plus Washington, D.C. had legalized cannabis use.

Prevalence of cannabis use has increased overall since 2007 which has raised concerns about potential negative consequences associated with problematic use, specifically cannabis use disorder. "These latest findings could directly inform policymakers and public health practitioners about the degree to which other broader contextual factors also influence cannabis use patterns in the U.S.," noted Philbin.

States were categorized as liberal, moderate, or conservative based on the 2005 and 2011 State Rank on Policy Liberalism Index, which is based on policy indicators for which liberals and conservatives commonly differ. The Index ranked each state from 1 (most liberal) to 50 (most conservative) based on its policies regulating gun control, abortion access, Temporary Assistance to Needy families, collective bargaining, and tax structure.

Average state-level prevalence of past-year cannabis use by age was lowest for ages 26 and older and highest for ages 18 to 25 throughout the study period. Average prevalence increased for ages 18-25 in liberal states, from 33 percent to 37 percent, and rose marginally in conservative states, from 25 percent to 26 percent. The same pattern of use was observed for ages 26 and over in liberal (8 percent to 10 percent) and conservative (6 percent to 7 percent) states. For ages 12-17, however, past year use did not significantly change from 2004-2006 to

2010-2012 in liberal or conservative states.

In contrast, cannabis use disorder among past-year cannabis users decreased from 2004-2006 to 2010-2012 among those aged 18-25 in conservative states (22 percent to 18 percent) and liberal states (20 percent to 17 percent). Among individuals ages 26 and over, cannabis use disorder among past-year users decreased in liberal states (11 percent to 8 percent). For 12-17 year olds, cannabis use disorder decreased in conservative states (28 percent to 25 percent), though still remained marginally higher than in liberal [states](#) (24 percent).

"Our study highlights the need for researchers and public health professionals to distinguish between cannabis use and cannabis use disorder when interacting with patients at the individual level and when developing primary prevention strategies and interventions at the population level," said Silvia Martins, MD, Ph.D., associate professor of Epidemiology and senior author. "This line of research not only helps identify how state-level policies as a whole impact cannabis use outcomes, but ultimately supports the development of more health-promoting policies."

**More information:** *International Journal of Drug Policy* (2019). [DOI: 10.1016/j.drugpo.2018.10.010](https://doi.org/10.1016/j.drugpo.2018.10.010)

Provided by Columbia University's Mailman School of Public Health

Citation: Cannabis use disorder: The policy climate matters (2019, January 23) retrieved 27 April 2024 from <https://medicalxpress.com/news/2019-01-cannabis-disorder-policy-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.