


CDC: Proportion of increased-risk deceased organ donors on rise

January 25 2019



(HealthDay)—Among deceased organ donors, there has been an increase



in the proportion at increased risk for transmitting hepatitis B virus (HBV), hepatitis C virus (HCV), and HIV to recipients, according to research published in the Jan. 25 issue of the U.S. Centers for Disease Control and Prevention *Morbidity and Mortality Weekly Report*.

Winston E. Abara, M.D., from the CDC in Atlanta, and colleagues analyzed deceased donor data for 2010 to 2017 reported to the Organ Procurement and Transplantation Network for increased-risk donors (IRDs) and standard-risk donors (SRDs).

The researchers noted an approximate 200 percent increase in the proportion of IRDs during the period, from 8.9 to 26.3 percent. There was also an approximate 200 percent increase in the percentage of deceased donors with drug intoxication as the mechanism of death, from 4.3 to 13.4 percent. About a 500 percent increase was seen in the proportion of donors with reported injection drug use, from 1.3 to 8.0 percent. IRDs were significantly more likely to have positive HBV and HCV screening results compared with SRDs.

"Identification of risk factors for viral bloodborne pathogen infection among <u>organ donors</u> is nonetheless important so that recipients and their clinicians can be notified and patients can be appropriately screened posttransplant," the authors write.

More information: Abstract/Full Text

Copyright © 2019 HealthDay. All rights reserved.

Citation: CDC: Proportion of increased-risk deceased organ donors on rise (2019, January 25) retrieved 17 April 2024 from

https://medicalxpress.com/news/2019-01-cdc-proportion-increased-risk-deceased-donors.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.