

## **CDC: Slight hike in prevalence of gastroschisis since 2006-2010**

January 23 2019

---



(HealthDay)—Over time, the prevalence of gastroschisis has increased,

with more babies born with gastroschisis in areas with high and medium versus low opioid prescription rates, according to research published in the Jan. 18 issue of the U.S. Centers for Disease Control and Prevention *Morbidity and Mortality Weekly Report*.

Tyiesha D. Short, M.P.H., from the CDC in Atlanta, and colleagues pooled data from 20 population-based state surveillance programs and examined age-specific gastroschisis prevalence during 2006 to 2010 and 2011 to 2015. Annual gastroschisis prevalence was compared by annual opioid prescription rate categories using an ecologic approach.

The researchers found that the prevalence of gastroschisis increased 10 percent from 2006-2010 to 2011-2015 (prevalence ratio, 1.10; 95 percent confidence interval, 1.0 to 1.1); prevalence was highest among mothers aged

Citation: CDC: Slight hike in prevalence of gastroschisis since 2006-2010 (2019, January 23) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2019-01-cdc-slight-hike-prevalence-gastroschisis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.