

# Cigarette smoking associated with increased risk of peripheral artery disease in African-Americans

January 23 2019

---



Credit: CC0 Public Domain

African Americans who smoke cigarettes are more likely than those who don't smoke to develop peripheral artery disease, according to new research in *Journal of the American Heart Association*, the Open Access

Journal of the American Heart Association/American Stroke Association.

Peripheral artery disease (PAD) is a narrowing of [arteries](#) other than those directly serving the heart. These [blood vessels](#) provide [blood flow](#) to the arms, legs, the brain and other organs such as the kidneys. If this blood flow is reduced, stroke, kidney failure, erectile dysfunction, pain in the legs when walking and even loss of limbs can occur. PAD is associated with several heart disease risk factors including hypertension, diabetes and obesity.

African Americans are disproportionately affected by PAD, nearly twice as much as non-Hispanic Whites. While cigarette smoking is a recognized PAD risk factor, little is known about its specific impact on African Americans.

"Peripheral artery disease is a major disease burden in the African American population," said Donald Clark III, M.D., M.P.H., assistant professor of medicine at the University of Mississippi Medical Center in Jackson. "Establishing a clear association between smoking and peripheral artery disease in this particularly vulnerable population can help guide our efforts to lower their risk and to help them improve their health."

To learn more, researchers analyzed [medical records](#) of about 5,300 participants, aged 21 to 84 years, enrolled in the Jackson Heart Study, the largest single-site cohort study investigating cardiovascular risk factors in African Americans. Among the participants who self-reported, smoking status was 68 percent who never smoked (3579 individuals), 19 percent (986) past smokers and 13 percent (693) current smokers.

To determine evidence of PAD, the researchers looked at measures of [peripheral artery disease](#), including: ankle-brachial index, which

compares blood pressure in the arteries of the extremities, and computer tomography (CT) scanning to identify calcium build-up in blood vessels, which can indicate the presence of artery-blocking plaques.

In addition to being twice as likely as non-smokers to have subclinical PAD in their lower extremities, current smokers were eight times more likely to have high calcium build-up in the aorta, the body's main artery.

The researchers also found an association between the number of cigarettes smoked and the likelihood of PAD. Participants who smoked 20 or more cigarettes a day had significantly higher odds of PAD compared to those who smoked less than 20 cigarettes each day.

The study was conducted through the support of the American Heart Association's Tobacco Regulation and Addiction Center, as part of a multi-year grant initiative from the U.S. Food and Drug Administration (FDA) and the National Institutes of Health (NIH). The aim of the initiative is to conduct research to document the adverse cardiovascular effects of tobacco products that can inform the FDA's regulation of these products, to learn more about the attitudes and behaviors associated with tobacco use and to help shape effective communication about the health effects of tobacco products. The Association's Center has focused on vulnerable populations in an effort to promote health equity.

"The findings from this study give us strong evidence of the specific debilitating and life-threatening risks African American smokers face, especially the more they smoke," said Mariell Jessup, M.D., Chief Science and Medical Officer of the American Heart Association. "This type of research can be useful in the development of clear messages targeted to our African American population to underscore the real physical costs of tobacco product use."

**More information:** Donald Clark et al, Cigarette Smoking and Subclinical Peripheral Arterial Disease in Blacks of the Jackson Heart Study, *Journal of the American Heart Association* (2019). DOI: 10.1161/JAHA.118.010674

Provided by American Heart Association

Citation: Cigarette smoking associated with increased risk of peripheral artery disease in African-Americans (2019, January 23) retrieved 8 May 2024 from <https://medicalxpress.com/news/2019-01-cigarette-peripheral-artery-disease-african-americans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.