

Depression and obesity linked to greater likelihood of hip pain

January 9 2019




Illustration of Hip (Frontal view). Credit: Wikipedia

In a representative sample of the German population, older age, obesity, and depressive disorder were associated with experiencing chronic hip pain.

In the *Musculoskeletal Care* study of 2,515 adults, 4.9 percent reported

chronic hip pain and an additional 1.5 percent reported chronic hip pain that was disabling. Hip pain affected 1-5 sites in 47 percent, and was widespread (6-19 sites) in 50 percent.

Obesity and increased values on a depression scale were associated with a 2.55-times and 8.53-times higher likelihood of chronic hip pain, respectively, compared with individuals without pain. Increased values on the depression scale also increased the likelihood of experiencing disabling chronic hip pain.

More information: *Musculoskeletal Care* DOI: [10.1002/msc.1380](https://doi.org/10.1002/msc.1380)

Provided by Wiley

Citation: Depression and obesity linked to greater likelihood of hip pain (2019, January 9) retrieved 17 May 2024 from <https://medicalxpress.com/news/2019-01-depression-obesity-linked-greater-likelihood.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
