


## Stroke produces dysfunctional brain cells

January 7 2019



Uncoupled intrinsic and synaptic excitability in DCX+ ABGCs after stroke. Credit: Ceanga et al., *JNeurosci* (2019)

Mice produce new neurons in the hippocampus following a stroke that fail to develop properly, finds new research published in *JNeurosci*. Intervening in the production of these cells may help to mitigate stroke-induced memory impairments.

Stroke has long been known to increase adult neurogenesis. Despite the



proliferation of new cells in a brain region critical for memory, previous stroke research in animals shows this process is accompanied by deficits on tasks that depend on the hippocampus. These observations led Albrecht Kunze and colleagues to investigate how newborn cells mature and integrate into the existing hippocampal network after stroke.

By temporarily cutting off <u>blood supply</u> to the brains of male and female mice, the researchers demonstrate the neurons generated as a result of this stroke model develop into hyperexcitable cells that may contribute to hippocampal dysfunction.

This finding begins to uncover the cellular mechanisms underlying poststroke neuropsychiatric disorders.

**More information:** Stroke accelerates and uncouples intrinsic and synaptic excitability maturation of mouse hippocampal DCX+ adult-born granule cells, *JNeurosci* (2019). <u>DOI:</u> 10.1523/JNEUROSCI.3303-17.2018

## Provided by Society for Neuroscience

Citation: Stroke produces dysfunctional brain cells (2019, January 7) retrieved 3 May 2024 from <a href="https://medicalxpress.com/news/2019-01-dysfunctional-brain-cells.html">https://medicalxpress.com/news/2019-01-dysfunctional-brain-cells.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.