

How does ecstasy kill?

January 23 2019, by Nicole Lee



Credit: Ann H from Pexels

MDMA (Methylenedioxymethamphetamine), commonly referred to as ecstasy, was manufactured as a potential pharmaceutical early last century. It had some limited use in the 1970s as a therapeutic aid in trauma treatment and in relationship counselling, and more [recent studies](#) using MDMA for trauma have shown some promise.

Structurally, MDMA is similar to the stimulant methamphetamine and to the hallucinogen mescaline, and so has both stimulant and mildly hallucinogenic effects.

Most problems with recreational MDMA are acute. Dependence and other long-term problems are quite rare. Less than 1% of all [drug treatment presentations](#) are for ongoing problems with MDMA, such as dependence.

Most fatalities from taking ecstasy are a result of a [combination of factors](#), not just the drug itself.

Most of these conditions don't result in death if they are treated early, but because of the stigma associated with using [illicit drugs](#), sometimes people don't seek help early enough. Any unusual or unwanted symptoms experienced while taking ecstasy should be treated as soon as they appear.

Contaminants and polydrug use

Most people are under the impression drugs are illegal because they are dangerous, but a [drug's legal status](#) isn't necessarily related to relative danger. In fact, drugs are much more dangerous because they are unregulated, manufactured by backyard chemists in clandestine laboratories.

Unlike alcohol, which is a highly regulated drug, there's no way to tell how potent illicit drugs are or what's in them, unless you test them.

In Australia, what is sold as ecstasy may contain a lot of MDMA or very little. Pills can contain other more dangerous drugs that mimic the effects of MDMA, and benign substances, such as lactose, as filler agents.

A recent report on findings from Australia's first official pill testing trial at the Groovin' the Moo music festival last year, found nearly [half the pills](#) tested were of low purity. Some 84% of people who had their pills tested thought they had bought MDMA but only 51% actually contained any MDMA.

Some of the more dangerous contaminants found in pills include [PMA](#) (paramethoxyamphetamine), which is more toxic at lower doses than ecstasy; [N-Ethylpentylone](#), a [cathinone](#) which is a lot more potent than MDMA making it easier to take too much; and [NBOMes](#) (N-methoxybenzyl), which is more toxic at lower doses than other hallucinogenic drugs and can cause [heart attack](#), [renal failure](#), and stroke.

Pills have also been detected in UK and NZ with up to [three doses of MDMA](#) in a single pill.

Although it's possible to take too much MDMA and experience severe toxic effects, as with other illicit drugs, most [ecstasy-related deaths](#) involve multiple drugs.

Sometimes these drug mixes are unexpected and sometimes people take multiple drugs deliberately. It's safer for people using ecstasy to limit use of other drugs, including alcohol, to avoid risk of adverse effects.

Heatstroke

Heatstroke or hyperthermia (dangerously high body temperature) is one of the most common issues among people taking MDMA.

MDMA increases [body temperature](#) and sweating, and using it is often accompanied by physical activity (such as dancing) in a hot environment (such as a crowded venue or in the summer heat), exacerbating fluid loss. If you don't have enough fluids your body can't cool itself properly.

The effect of ecstasy can be exacerbated by consuming alcohol. Alcohol is a diuretic, so it makes you urinate more and increases dehydration. Dehydration [increases risk](#) of heatstroke.



Credit: AI-generated image ([disclaimer](#))

Heatstroke can cause brain, heart, kidney and muscle damage, and if left untreated can cause serious complications or death.

If active, people taking MDMA should drink around 500ml (two cups) of water an hour and take regular breaks. Isotonic drinks (such as Powerade and Gatorade) are also OK.

Water intoxication

People using MDMA can get really thirsty. Some is probably the direct effect of MDMA, some because they're hot, and some from dehydration.

But if you have too much water the ratio of salts and water in the body becomes unbalanced – basically the level of salt in your body gets too low and your cells start swelling with water. The technical name is [hyponatraemia](#).

MDMA is an anti-diuretic, so it makes you retain water, which can [increase risk](#) of water intoxication.

People may feel nausea with vomiting, confusion, severe fatigue, muscle weakness and cramps.

People taking ecstasy need to stay hydrated but only replace what is lost through sweating – around 500ml per hour if active and around 250ml an hour when inactive.

Serotonin syndrome

The main action of MDMA in the brain is an increase in [serotonin](#), which among other things is responsible for regulating pro-social behaviour, empathy and optimism. This is why people who have taken MDMA feel connection with and positivity towards others.

But too much serotonin can result in "[serotonin syndrome](#)". It typically occurs when other drugs that also raise serotonin levels (other stimulants, antidepressants) are taken together with MDMA.

Signs include high body temperature, agitation, confusion, problems controlling muscles, headache and the shakes. People might also experience seizures or loss of consciousness.

It can be fatal if the symptoms are left untreated, so if anyone taking MDMA shows any of these signs they should be treated immediately. It's safer not to mix different types of drugs, especially if you do not know what's in them.

Other causes

More rarely, fatalities have been reported as a result of other health complications after taking [ecstasy](#), especially if the person has pre-existing risk factors, such as high blood pressure or a heart condition. Complications related to heart failure, liver failure and brain haemorrhage have been reported in people already at high risk of these problems.

The number of people who die from party drugs is [relatively low](#) compared to other drugs such as heroin, alcohol, and pharmaceuticals. But the media tend to [report](#) a higher proportion of these deaths compared to other drugs, increasing the perception of harm. Most of the deaths are not directly from the drug itself but other complications or contaminants.

It's safest not to take drugs at all, but if you choose to, it's safer to take a small amount first (like a quarter of a pill) and wait at least an hour to make sure there are no ill effects; drink about 500ml per hour of water if active; and don't mix drugs, including alcohol.

In the absence of a legal, uncontaminated supply of MDMA, when pill testing becomes available in Australia it will at least help people make informed decisions about drug use and [reduce the risk](#) of fatalities and other harms. People often [choose not to take](#) their pills, or take smaller amounts, when they discover contaminants.

This article is republished from [The Conversation](#) under a Creative

Commons license. Read the [original article](#).

Provided by The Conversation

Citation: How does ecstasy kill? (2019, January 23) retrieved 29 April 2024 from <https://medicalxpress.com/news/2019-01-ecstasy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.