

Emergency/urgent hospitalizations linked to accelerated cognitive decline in older adults

January 12 2019



Credit: CC0 Public Domain

Emergency and urgent hospitalizations are associated with an increased rate of cognitive decline in older adults, report researchers at Rush University Medical Center. Results of their study, published in the Jan.

11, 2019, online issue of *Neurology*, the medical journal of the American Academy of Neurology, shows that hospitalization may be a more of a major risk factor for long-term cognitive decline in older adults than previously recognized.

"We found that those who have non-elective (emergency or urgent) hospitalizations and who have not previously been diagnosed with dementia or Alzheimer's disease had a [rapid decline](#) in cognitive function (i.e., thinking abilities) compared to the prehospital rates," said Bryan James, Ph.D., an epidemiologist and in the Rush Alzheimer's Disease Center and an assistant professor in the Rush Department of Internal Medicine. "By comparison, people who were never hospitalized and those who had elective hospitalizations did not experience the drastic decline in cognitive function."

In 2017, James and colleagues presented a preliminary version of their study results at the Alzheimer's Association International Conference in London.

Study compares hospitalization data and cognitive assessments for 777 older adults

The data emerged from a study of 777 older adults (81 years old on average, 75 percent of them women) enrolled in the Rush Memory and Aging Project (MAP) in Chicago. The study involved annual cognitive assessments and clinical evaluations.

Information on hospitalizations was acquired by linking records of 1999 to 2010 Medicare claims for these participants with their MAP data. All hospital admissions were designated as elective, emergency, or urgent. The latter two were combined as non-elective for analysis.

Of the 777 participants, 460 were hospitalized at least once over an average of almost five years of observation. Of those who were hospitalized, 222 (29 percent of the total study population) had at least one elective hospital admission, and 418 (54 percent) had at least one non-elective hospital admission. These groups included 180 participants (23 percent) who had both types of hospitalizations.

Non-elective hospitalizations were associated with an approximately 50 percent acceleration in the rate of cognitive decline from before hospitalization, and a rate of cognitive decline that was more than double the rate in persons who were not hospitalized. Elective hospitalizations, however, were not associated with acceleration in the rate of decline at all.

'Elective admissions do not necessarily carry the same risk'

"We saw a clear distinction: non-elective admissions drive the association between hospitalization and long-term changes in cognitive function in later life, while elective admissions do not necessarily carry the same risk of negative cognitive outcomes," James said. "These findings have important implications for the medical decision making and care of older adults.

"While recognizing that all medical procedures carry some degree of risk, this study implies that planned hospital encounters may not be as dangerous to the cognitive health of older persons as emergency or urgent situations."

It is unknown why emergency and urgent hospitalizations carry a higher risk of long-term cognitive decline than elective hospitalizations, but it could be due to differences in levels of sickness (though the authors

controlled for health status), stress, or [hospital](#) procedures involved. The authors plan to explore these reasons in future research.

This work expands upon previous research which has shown that after being hospitalized, older adults are at high risk for memory and other cognitive problems, including both transient (temporary) delirium and long-term changes in cognition, including dementia. According to the Healthcare Cost and Utilization Project in October 2010, 40 percent of all hospitalized patients in U.S. are age 65 and older. Therefore, hospitalization may be an under-recognized risk factor for cognitive decline and dementia for a large number of older adults that deserves more attention.

Detection of dementia at the earliest stages has become a worldwide priority, because drug treatments, prevention strategies and other interventions will likely be more effective very early in the disease process, before extensive brain damage has occurred.

Provided by Rush University Medical Center

Citation: Emergency/urgent hospitalizations linked to accelerated cognitive decline in older adults (2019, January 12) retrieved 10 April 2024 from <https://medicalxpress.com/news/2019-01-emergencyurgent-hospitalizations-linked-cognitive-decline.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
