

Study: Faster weight loss no better than slow weight loss for health benefits

January 29 2019



Credit: CC0 Public Domain

Losing weight slowly or quickly won't tip the scale in your favour when it comes to overall health, according to new research. Health researchers at York University found that people who lose weight quickly versus

those who lose it slowly don't get any additional health benefits and it's the amount of weight lost overall that can have an impact.

In the study led by Jennifer Kuk, associate professor in York University's Faculty of Health, researchers looked at the data of over 11,000 patients at a publicly-funded clinical [weight](#) management program and found that those who lost weight quickly had similar improvements in metabolic health with those who lost weight slowly. Moreover, the rate of weight loss matters less for [overall health](#) benefits than the amount of weight you lose.

Normally, individuals are recommended to lose weight at one to two pounds per week, as faster weight loss is related with a slightly [higher risk](#) for gallstones. However, there are reasons to believe that faster weight loss may have better effects for [cardiovascular disease](#) and diabetes risk factors.

The study is the first study of its kind to look specifically at risk factors for cardiovascular health and diabetes.

"With the same pound for pound weight loss, there is no difference in terms of health benefits if you lose weight fast or slow," says Kuk.

"However, given the risk for gallstones with faster weight loss, trying to lose weight at the recommended one to two pounds per week is the safer option."

The study looked at 11,283 patients who attended the Wharton Medical Clinic Weight Management Program between July 2008 and July 2017. Researchers found that patients who lost weight more quickly tended to have a bigger reduction in obesity and better health improvements than patients who lost weight slowly. However, these improvements in [health](#) associated with faster weight loss were abolished after adjusting for absolute weight loss.

"The results show that we really need to look at interventions that focus on long-term weight management that can achieve sustained [weight loss](#) at the recommended one to two pounds per week," says Kuk.

The study is published today in the *Journal of Obesity*.

More information: *Journal of Obesity* (2019). [DOI: 10.1155/2019/3609642](#) ,
www.hindawi.com/journals/job/2019/3609642/

Provided by York University

Citation: Study: Faster weight loss no better than slow weight loss for health benefits (2019, January 29) retrieved 7 May 2024 from <https://medicalxpress.com/news/2019-01-faster-weight-loss-health-benefits.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.