

Heart patients prescribed opioids at hospital discharge often fare worse

January 31 2019



Credit: American Heart Association

Heart disease patients prescribed opioids at hospital discharge were less likely to follow up with their healthcare provider or to participate in heart rehabilitation than patients who were not, according to new research in *Journal of the American Heart Association*, the Open Access

Journal of the American Heart Association/American Stroke Association.

As the nationwide opioid epidemic continues, the risks versus benefits of these strong painkillers are being questioned. Previous research found [health problems](#) among [patients](#) prescribed opioids for non-cancer-related pain and a recent study linked opioid use to an increase risk of atrial fibrillation.

In this study, researchers analyzed 2,495 patients (mostly white males; average age 60) discharged from Vanderbilt University Medical Center for heart attack, sudden heart failure, or both between October 2011 and December 2015. Twenty percent of patients were discharged with an opioid prescription.

Compared to those not prescribed opioids at discharge, researchers found patients prescribed opioids were:

- less likely to follow up with their healthcare provider or participate in heart rehabilitation thirty days after discharge; and
- slightly more likely to visit the [emergency room](#), be readmitted to the hospital, or to die within 90 days.

"Hospital discharge provides a unique opportunity to evaluate each patient's medication regimen," said study lead author Justin S. Liberman, M.D., M.P.H., assistant professor of anesthesiology at Vanderbilt University Medical Center in Nashville, Tennessee. "It's important for [healthcare providers](#) to understand the other ways that opioids may affect a patient's future interactions with the healthcare system."

Researchers reviewed patients' medical records through the Vanderbilt Inpatient Cohort Study. After [hospital discharge](#), participants provided information by telephone interview about their use of medical services.

Interviews occurred 2-3 days, 30 days and 90 days after discharge.

The study shows an association but does not prove that [opioid](#) prescriptions caused patients to miss follow-up care. Researchers said, it's possible that diminished physical and mental function, known effects of opioids, contributed.

The study may be limited by a lack of information about the number of opioids taken and their duration of use. Also, the study focused on one hospital with a primarily white population and may not apply to others.

More information: Justin S. Liberman et al. Opioid Prescriptions at Hospital Discharge Are Associated With More Postdischarge Healthcare Utilization, *Journal of the American Heart Association* (2019). [DOI: 10.1161/JAHA.118.010664](#)

Provided by American Heart Association

Citation: Heart patients prescribed opioids at hospital discharge often fare worse (2019, January 31) retrieved 20 April 2024 from <https://medicalxpress.com/news/2019-01-heart-patients-opioids-hospital-discharge.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--