


Prescribed opioids raise risk of pneumonia in patients with and without HIV

January 7 2019



A black and white X-ray picture showing a triangular white area on the left side. A circle highlights the area. Credit: James Heilman, MD./Wikipedia

Taking prescribed opioids raises the risk of pneumonia in individuals with and without HIV, a new Yale-led study finds.

The study, published in *JAMA Internal Medicine*, reinforces the concern that prescription opioid painkillers have a [negative impact](#) on the [immune system](#). It also underscores the need for awareness among opioid prescribers who can take steps to minimize the risk of [pneumonia](#) through vaccination and promoting smoking cessation, the researchers said. To their knowledge, this is the first study to examine the impact of prescribed opioids on pneumonia risk in patients with HIV.

Opioids are commonly prescribed to individuals for pain, especially those with HIV. However, there's evidence that some opioids—including codeine, fentanyl, and morphine—suppress the immune system and limit the body's ability to fight bacterial infections, such as pneumonia. To investigate the link between prescribed opioids and pneumonia, the research team analyzed data from patients enrolled in the Veterans Aging Cohort Study, a national study of individuals who receive care through the Veterans Health Administration (VA).

The [research team](#) conducted the study using data from [individuals](#) who were treated at a VA between 2000 and 2012. They included patients living with and without HIV. The team matched patients hospitalized for pneumonia with similar patients who did not have pneumonia. In their analysis, the researchers looked at length of time patients were taking opioids, as well as dosages and whether the medications had known immunosuppressive properties or not.

The researchers found that patients prescribed medium or high doses of opioid painkillers were at greater risk for pneumonia than those not taking the medications. The risk was compounded if the [opioid](#) contained immunosuppressive properties. Individuals with HIV tended to be more likely to develop pneumonia even at low doses of opioids and

especially with immunosuppressive opioids.

"We saw that prescription opioids were independently associated with pneumonia requiring hospitalization," said E. Jennifer Edelman, M.D., the corresponding author and an associate professor at Yale School of Medicine.

Prescription opioids can impact the body's defenses against pneumonia in different ways, the researchers said, including by suppressing cough, respiration, and mucus secretion. The study "lends credence to the hypothesis that opioids have effects on the immune system that are clinically relevant," Edelman noted.

In addition to raising awareness of the increased pneumonia risk among providers and patients, the findings could lead to changes to mitigate that risk and add to the growing list of potential harms associated with these medications. Providers could consider prescribing lower doses of opioids or opioids that do not suppress the immune system, the researchers said. They can also be proactive about getting [patients](#), especially those with HIV, vaccinated against pneumonia.

Provided by Yale University

Citation: Prescribed opioids raise risk of pneumonia in patients with and without HIV (2019, January 7) retrieved 18 April 2024 from <https://medicalxpress.com/news/2019-01-opioids-pneumonia-patients-hiv.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.