

Persistent low body weight in young kids increases risk for anorexia nervosa later

January 31 2019



Study first author Zeynep Yilmaz, PhD. Credit: UNC School of Medicine

A new study has found that a persistent low body mass index (BMI) in children, starting as young as age 2 for boys and 4 for girls, may be a risk factor for the development of anorexia nervosa in adolescence.

In addition, the study, published in the February 2019 issue of the *Journal of the American Academy of Child and Adolescent Psychiatry*, found that a persistent high BMI in childhood may be a risk factor for later development of bulimia nervosa, binge-eating disorder, and purging disorder. This large population study is based on analysis of data from 1,502 individuals who participated in the Avon Longitudinal Study of Parents and Children in the UK.

"Until now, we have had very little guidance on how to identify children who might be at increased risk for developing eating disorders later in adolescence," said Zeynep Yilmaz, Ph.D., study first author and an assistant professor of psychiatry and genetics at the UNC Center of Excellence for Eating Disorders in the University of North Carolina School of Medicine. "By looking at growth records of thousands of children across time, we saw early warning profiles that could signal children at risk."

Co-author Cynthia Bulik, Ph.D., Distinguished Professor of Eating Disorders also from UNC highlights, "Clinically, this means that pediatricians should be alert for children who fall off and stay below the growth curve throughout childhood. This could be an early warning sign of risk for anorexia nervosa. The same holds for [children](#) who exceed and remain above the growth curve—only their risk is increased for the other eating disorders such as bulimia nervosa and binge-eating disorder."

Yilmaz said that although eating disorders are psychiatric in nature, the study highlights the need to also consider metabolic [risk factors](#) alongside psychological, sociocultural, and environmental components.

"The differences in childhood body weight of adolescents who later developed eating [disorders](#) started to emerge at a very early age—way too early to be caused by social pressures to be thin or dieting. A more likely explanation is that underlying metabolic factors that are driven by genetics, could predispose these individuals to weight dysregulation. This aligns with our other genetic work that has highlighted a metabolic component to anorexia nervosa."

Corresponding author of the study is Nadia Micali, MD, MRCPsych Ph.D., Full Professor at University of Geneva Faculty of Medicine and Head of Geneva University Hospitals' Division of Child and Adolescent Psychiatry.

"Our results also highlight the multi-factorial composition of [eating disorders](#), as well as the need to develop early detection tools that could be used as part of routine checks by all pediatricians. Indeed, the earlier the problem is identified, the better it can be managed, especially if support is provided to the family as a whole, rather than just the individual," Micali said.

More information: Zeynep Yilmaz et al, Developmental Premorbid Body Mass Index Trajectories of Adolescents With Eating Disorders in a Longitudinal Population Cohort, *Journal of the American Academy of Child & Adolescent Psychiatry* (2018). [DOI: 10.1016/j.jaac.2018.11.008](https://doi.org/10.1016/j.jaac.2018.11.008)

Provided by University of North Carolina Health Care

Citation: Persistent low body weight in young kids increases risk for anorexia nervosa later (2019, January 31) retrieved 23 June 2024 from <https://medicalxpress.com/news/2019-01-persistent-body-weight-young-kids.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.