

## Short bouts of intense activity may have more benefits than hours of lower exertion

January 7 2019



Credit: CC0 Public Domain

The expectation of constant availability due to our 21st-century, technology-driven lifestyles can be exhausting. Throw in laundry,



shopping and social obligations, and finding the time to exercise can seem impossible.

In fact, the single most common excuse for not exercising regularly is a perceived lack of time, said Glenn Gaesser, professor of exercise and wellness at Arizona State University's College of Health Solutions.

He and ASU Assistant Professor Siddhartha Angadi have been working together for over a decade to research the effects of <u>high-intensity</u> <u>interval training</u>, or HIIT—characterized by short bursts of intense activity—on various health outcomes.

Their research reflects recent <u>updates</u> to the national Physical Activity Guidelines, a set of recommendations that impact public health policies.

Experts have long agreed that adults should aim for roughly 150 minutes of moderate physical activity per week. When the guidelines were first published in 2008, it was thought that in order to obtain meaningful health benefits, those 150 minutes had to be accrued in bouts of at least 30 minutes of activity at a time.

Gaesser and Angadi's research has shown that not only can shorter bouts of physical activity produce the same benefits as longer bouts, but that if the shorter bouts are ramped up from a moderate level (something akin to a brisk walk) to a vigorous level (where you're almost out of breath but not quite), they may even produce more health benefits than longer, moderate-level bouts.

For instance, in more than one study, Gaesser and Angadi found that HIIT was better than traditional, continuous bouts of exercise at lowering blood pressure.

"Short bouts always do as well and sometimes better than longer bouts,"



Gaesser said.

A common misconception about HIIT is that it's too difficult for the average person. In actuality, HIIT is tailored to an individual's personal capabilities. The principle is to do just enough physical activity to elevate your <u>heart rate</u> to the point just before you start to feel fatigued—which is different for everyone—and then stop. That's one HIIT bout.

Generally, one HIIT bout takes anywhere from 30 seconds to a few minutes. You can rest for as much or as little time between each bout as you like, as long as you do enough bouts each week to add up to 150 minutes. That makes it easy to incorporate HIIT into a busy schedule; on a regular workday, you can get 30 minutes of physical activity in by taking a few minutes' break every hour to climb the stairwell or take a brisk walk around the office building.

"People might not be able to do a 30-minute routine every day," Gaesser said, "but if you say you don't have two minutes here and there, I'm just not buying it."

Over the years, he and Angadi have seen everyone from obese individuals to <u>heart-failure patients</u> employ HIIT with ease. They're currently in the midst of a study looking at the effects of HIIT on the hearts of breast cancer patients.

Breast cancer is highly survivable, with about a 90 percent five-year survival rate. But in women who are over the age of 60, <u>heart disease</u> competes with the cancer as a cause of death because chemotherapy is so hard on the heart. It causes oxidative stress, so a natural reaction would be to dose patients with antioxidants like vitamins C and E. However, antioxidants go everywhere in the body, including into the cancer cells, which actually helps them fight the chemo and spread further.



"Because of this, there was a desperate need to come up with a strategy to protect the heart without compromising the cancer-killing properties of chemotherapy," said Angadi.

Based on his and Gaesser's research, he knew that HIIT builds up <u>antioxidants</u> inside tissues used while exercising—most notably in this case, the heart. So he reasoned that by "dosing" breast cancer patients with HIIT before they undergo chemotherapy, their hearts could be preconditioned to better withstand the damage it caused.

In an animal study, less than 10 percent of rats dosed with HIIT before receiving chemotherapy died from cardiac-related reasons, while roughly 50 percent of rats who were not dosed with HIIT before chemotherapy died from heart failure.

"It was night and day," said Angadi. "And when we looked at their hearts via electron microscopy, the hearts that got the chemo without exercise were just wrecked. Whereas the hearts that got exercise before the chemo were relatively preserved."

Initial results of the first human trials taking place at Angadi and Gaesser's lab in downtown Phoenix are promising, indicating that hearts of patients dosed with HIIT are faring better than those in a control group who were only asked to walk 10,000 steps each day.

While some cancer centers are now incorporating exercise into patients' treatment regimens, it is usually the case that it is only introduced after chemo has already begun. By then, Angadi said, it's too late.

"Chemotherapy causes direct cardiac tissue death, and these are specialized tissues in our body," he said. "Once they're gone, they're gone. There's no bringing them back. So you have to act before the house catches fire."



Depending on the success of the breast cancer patient study, Angadi and Gaesser hope to move onto studying the effects of HIIT in children with leukemia, for which the same cardiotoxic drugs are used in treatment. Like <u>breast cancer</u>, pediatric leukemia also has a high survival rate but can wreak havoc on the heart, causing survivors of the disease to experience cardiac events much earlier in life, usually in their early 40s.

"The drugs (they're given for treatment) are effectively <u>heart</u> disease and diabetes in a pill," Angadi said. "Don't get me wrong, you don't get the drugs, you die. You don't even make it to 40. So they're absolutely essential. But we need to come up with harm-reduction strategies to mitigate the risk associated with them."

Provided by Arizona State University

Citation: Short bouts of intense activity may have more benefits than hours of lower exertion (2019, January 7) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2019-01-short-bouts-intense-benefits-hours.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.