

# Adhesion protein optimizes border

February 18 2019, by Leigh Macmillan

---



Credit: AI-generated image ([disclaimer](#))

The epithelial cells that line the intestines build a specialized cell surface—the "brush border"—that processes and absorbs nutrients, and defends against pathogens. The brush border consists of thousands of finger-like membrane protrusions (microvilli) on each cell.

Matthew Tyska, Ph.D., and colleagues previously implicated the adhesion protein CDHR2 in the tight packing of microvilli on cultured

cells. Now, they have generated mice lacking CDHR2 expression in the [intestinal tract](#) to explore the protein's role in building brush borders in vivo.

They reported in *Molecular Biology of the Cell* that mice missing intestinal CDHR2 were smaller than control mice. Examination of the intestinal lining revealed that intestinal microvilli were significantly shorter, had disordered packing and had a 30 percent reduction in packing density. The structural changes were linked to reduced levels of key enzymes and transporters in the brush border, which potentially explains the reduced body weight.

The findings show that CDHR2 elongates and maximizes intestinal microvilli for peak functional capacity.

**More information:** Julia A. Pinette et al. Brush border protocadherin CDHR2 promotes the elongation and maximized packing of microvilli in vivo, *Molecular Biology of the Cell* (2018). [DOI: 10.1091/mbc.E18-09-0558](#)

Provided by Vanderbilt University

Citation: Adhesion protein optimizes border (2019, February 18) retrieved 17 May 2024 from <https://medicalxpress.com/news/2019-02-adhesion-protein-optimizes-border.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--