


# Is alcohol consumption more helpful than harmful? It depends on your age

February 28 2019

---



Credit: CC0 Public Domain

Studies of health effects of alcohol consumption may underestimate the risks of imbibing, particularly for younger people, according to a new study in the *Journal of Studies on Alcohol and Drugs*.

The research demonstrates that although cohort studies—the type of observational investigations typically used to study health benefits and risks—sometimes show benefits from [moderate alcohol consumption](#), they tend to enroll people age 50 and older. Studying people at this age range eliminates all those who have died before age 50 because of alcohol consumption.

Simply put, "deceased persons cannot be enrolled in cohort studies," write the authors, led by Timothy Naimi, M.D., M.P.H., of the Boston Medical Center. "Those who are established drinkers at age 50 are 'survivors' of their alcohol consumption who [initially] might have been healthier or have had safer [drinking](#) patterns" compared with others.

In fact, more than 40% of overall deaths from alcohol occur before age 50, according to the study. This suggests that most current studies underestimate alcohol-related risk compared with what would be observed across the full age spectrum. Further, the study demonstrates that most study participants are not representative of all persons who begin to drink alcohol.

To obtain data for the current research, Naimi and his colleagues used the Alcohol-Related Disease Impact software application from the U.S. Centers for Disease Control and Prevention. The data used were derived from government statistics on cause of [death](#) and health benefits from 2006 to 2010 in the United States.

Naimi and colleagues found age was a large factor in deaths prevented by alcohol consumption. Some 35.8% of the total deaths caused by alcohol occurred in people ages 20 to 49. However, of the deaths determined to have been prevented by alcohol consumption, only 4.5% occurred in this younger age group.

Although those ages 65 and older saw a similar 35.0% of the mortality

caused by drinking, 80% of the deaths that were prevented by alcohol consumption occurred in this group.

A somewhat similar pattern emerged when the researchers looked at years of potential life lost as a result of drinking. Of the overall years of life lost, 58.4% occurred in those ages 20 to 49. But this younger group saw only 14.5% of the years of life saved from drinking.

People 65 and older accounted for 15% of the overall years of life lost from alcohol [consumption](#). However, more than 50% of the years of life saved occurred within this older group.

In general, their results show that [younger people](#) "are more likely to die from [alcohol consumption](#) than they are to die from a lack of drinking," according to the authors, whereas [older people](#) are the ones most likely to see the [health benefits](#) of moderate drinking often mentioned in the news.

"This study adds to the literature questioning protective effects for [alcohol](#) on all-cause mortality," the authors conclude. Nonetheless, they write that "most who choose to drink can do so with relatively low risk."

**More information:** Naimi, T. S., Stadtmueller, L. A., Chikritzhs, T., Stockwell, T., Zhao, J., Britton, A., Saitz, R., & Sherik, A. (2019). Alcohol, age, and mortality: Estimating selection bias due to premature death. *Journal of Studies on Alcohol and Drugs*, 80, 63-98. [DOI: 10.15288/jsad.2019.80.63](#)

Provided by Journal of Studies on Alcohol and Drugs

Citation: Is alcohol consumption more helpful than harmful? It depends on your age (2019,

February 28) retrieved 23 April 2024 from <https://medicalxpress.com/news/2019-02-alcohol-consumption-age.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.