

New study shows annual noninvasive stool test effective for colon cancer screening

February 25 2019



Thomas Imperiale, MD of Regenstrief Institute and Indiana University School of Medicine, a health services research scientist and a gastroenterologist studies screening for colorectal cancer. Credit: Regenstrief Institute

A new study by Indiana University School of Medicine and the Regenstrief Institute provides the strongest evidence to date to support recommendations that average risk patients can safely opt for an annual, easy-to-use home stool test instead of a screening colonoscopy.

The researchers reviewed and analyzed the findings of 31 studies with a total of 120,255 participants. Each individual had a FIT (short for fecal immunochemical test), which identifies hidden blood in stool. FIT results were compared to the finding of a subsequent [screening colonoscopy](#) and were found to have high detection rates for colorectal cancer.

The study results are published online in advance of print in the March 6, 2019 issue of *Annals of Internal Medicine*.

"Our analysis finds that FIT is a good 'pre-screening' test for average risk, asymptomatic adults, saving them hassle and the U.S. healthcare system costs," said Thomas Imperiale, MD, the lead author, a gastroenterologist with IU School of Medicine and a research scientist with the Regenstrief Institute. "If annual FIT results remain negative, FIT buys you time until colonoscopy may be required, and it could be the case that a colonoscopy for screening may never be necessary or required."

Colonoscopy is considered to be the gold standard in the United States for colon cancer screening, and it is often performed in average-risk, asymptomatic adults 50 and older. But, in other countries, such as the United Kingdom, Canada, and Italy, colonoscopy is typically performed for average risk individuals with no symptoms only if FIT results are positive.

Imperiale said the research indicates a similar approach could be safely deployed in the United States and elsewhere, but only if there is a system to ensure annual testing.

"We need a system that is population-based rather than the current opportunistic one—with better laboratory-physician coordination, dedicated follow-up, and assurance that colonoscopy is performed if FIT

results are positive," he said. "The power of FIT is in its repeat application and in many cases, repeatedly negative results."

Of the more than 4,000 English-language studies on FIT that Imperiale and Rachel Gruber, a Regenstrief Institute research coordinator, reviewed, only the 31 studies that were analyzed included both FIT and colonoscopy results for each study participant. Approximately a third of the participants with positive FIT results in these 31 studies were found to have an advanced precancerous polyp or actual colon cancer.

The U.S. Preventive Services Task Force currently recommends screening for [colorectal cancer](#) using fecal occult blood testing—the category that includes FIT—sigmoidoscopy, [colonoscopy](#), and other tests from age 50 to age 75, without preferentially recommending one particular [screening test](#).

Provided by Regenstrief Institute

Citation: New study shows annual noninvasive stool test effective for colon cancer screening (2019, February 25) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-02-annual-noninvasive-stool-effective-colon.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--