

Why Asian horses don't get sick with the flu

February 7 2019



A typical view of a lake in Mongolia where horses and wild birds live in close proximity. This shared environment is likely to facilitate the exposure of horses to avian influenza viruses. Credit: Ariunbaatar Barkhasbaatar, Wildlife Conservation Society

Avian influenza viruses infect horses in Mongolia but do not cause large outbreaks of disease because they failed to acquire key genetic changes

to enable greater cross-species transmissibility, according to a study published February 7 in the open-access journal *PLOS Pathogens* by Pablo Murcia of the MRC-University of Glasgow Centre for Virus Research, and colleagues. Future studies aiming at understanding those blocks to cross-species transmission using avian and equine influenza viruses will provide insight on the mechanisms and determinants that underpin influenza emergence in mammals.

Viral diseases pose a constant threat to humans and animals. Occasionally, viruses establish in new hosts, sometimes with devastating consequences. It is not clear what allows a virus to infect and become transmissible in a new population, but ecology and evolution play an important part in this process. Influenza A viruses constitute the archetypical example of emerging viruses: their main natural reservoir is in [wild birds](#), but they have also established in humans, pigs and horses. To better understand how influenza A viruses circulate in nature and the ecological and evolutionary factors mediating viral emergence, Murcia and colleagues sequenced more than 20 avian influenza viruses collected from wild birds in Mongolia. By combining field studies, phylogenetics and experimental infections, the authors documented with unprecedented detail the stages that precede initial outbreaks during viral emergence in nature.

The results revealed that the avian influenza viruses are genetically related to a [virus](#) that caused an equine influenza [outbreak](#) in 1989. Experimental infections showed that most avian influenza viruses replicate in the respiratory tract of the horse without causing any tissue damage. Moreover, horses in Mongolia have been regularly exposed to them over a broad geographical area without causing clinically evident outbreaks. The findings suggest that equine infections by avian viruses able to replicate in horses are more common than originally thought, and that the failure to acquire key genetic changes is the main barrier to disease emergence. According to the authors, surveillance efforts should

be directed to monitor for the presence of horse-adaptive mutations in viruses derived from birds.

"New [influenza](#) viruses fail to emerge in horses despite abundant ecological opportunities, Murcia adds. "This is likely due to the inability of the viruses to acquire the necessary changes in their genome to adapt to horses."

More information: Zhu H, Damdinjav B, Gonzalez G, Patrono LV, Ramirez-Mendoza H, Amat JAR, et al. (2019) Absence of adaptive evolution is the main barrier against influenza emergence in horses in Asia despite frequent virus interspecies transmission from wild birds. *PLoS Pathog* 15(2): e1007531. doi.org/10.1371/journal.ppat.1007531

Provided by Public Library of Science

Citation: Why Asian horses don't get sick with the flu (2019, February 7) retrieved 24 April 2024 from <https://medicalxpress.com/news/2019-02-asian-horses-dont-sick-flu.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--