

Children exposed to air pollution at school may be at greater risk of overweight and obesity

February 1 2019



Credit: CC0 Public Domain

Exposure to air pollution, particularly at school, could be associated with a higher risk of overweight and obesity during childhood. This is the



conclusion of a study by the Barcelona Institute for Global Health (ISGlobal), an institution supported by "la Caixa", performed with 2,660 children between 7 and 10 years of age from 39 schools in Barcelona.

A few studies have associated exposure to air pollution with a higher risk of child obesity and overweight. However, these studies focused mainly on exposure at home, without considering the school environment.

The study, published in *Environment International* and performed under the BREATHE project, assessed for the first time the relation between obesity and overweight risk and <u>air pollution levels</u> at school and home, the two microenvironments where schoolchildren spend most of their time.

On one hand, the research team collected data on the children's weight and height and calculated their body mass index and obesity and overweight status. On the other hand, they used sensors in the schoolyard to measure the levels of outdoor pollution—<u>nitrogen dioxide</u> (NO₂), elemental carbon (EC), particle matter (PM2.5) and ultrafine particles -during one week in summer and another week in winter. They also estimated exposure levels to NO₂, NOx, PM2.5, PM10 and PMcoarse at the home address.

"We observed that children exposed to medium or high levels of air pollution at school—ultrafine particles, NO₂, PM2.5 and EC—had a higher risk of obesity and overweight as compared to those exposed to lower levels", concludes first author Jeroen de Bont, researcher at ISGlobal and IDIAP Jordi Gol. Exposure to high levels of PM10 at home was also associated with <u>higher risk</u> of obesity or overweight during childhood, although in this case the analysis was done with estimates of exposure levels.

"The study has however some limitations, which means that the results



are to be cautiously interpreted", says Martine Vrijheid, ISGlobal researcher and study coordinator. "Being a cross-sectional study, we only have data at one time-point, and we do not have enough data to establish the nature of the association. To draw more solid conclusions, we need new longitudinal studies that follow the study participants over time", she adds.

Regarding the mechanisms linking air pollution and overweight, some animal studies suggest that pollution can induce <u>oxidative stress</u>, <u>insulin</u> <u>resistance</u> and systemic inflammation, factors that are known to contribute to obesity", explains de Bont.

Most children were exposed to air <u>pollution</u> levels above those recommended by the World Health Organisation (WHO), both at school and home. Specifically, over 75% were exposed to PM2.5 levels above those recommended ($10\mu g/m3$) and more than 50% breathed NO₂ levels above those considered as safe ($40\mu g/m3$).

More information: Jeroen de Bont et al, Ambient air pollution and overweight and obesity in school-aged children in Barcelona, Spain, *Environment International* (2019). DOI: 10.1016/j.envint.2019.01.048

Provided by Barcelona Institute for Global Health

Citation: Children exposed to air pollution at school may be at greater risk of overweight and obesity (2019, February 1) retrieved 9 May 2024 from https://medicalxpress.com/news/2019-02-children-exposed-air-pollution-school.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.