

# New study seeks to guide clinical treatment for older patients with aortic stenosis

February 4 2019

---



Credit: CC0 Public Domain

Why do some patients recover quickly after surgery, while others don't? That is an important question when treating older frail patients suffering from aortic stenosis. Lead author Dae Hyun Kim, M.D., M.P.H., Sc.D.,

and principle investigator Director Lewis A. Lipsitz in the Marcus Institute for Aging Research at Hebrew SeniorLife explore this question in a paper published today in the *Journal of the American Medical Association Internal Medicine*.

Aortic stenosis is a disease that more often than not develops late in life. It occurs in 2% of people over 65 and affects more men than women. The condition causes the aorta -the main artery that carries blood to the rest of the body—to constrict reducing blood flow from the heart. The disease poses a serious health threat to [older adults](#), impacting function and quality of life at best, and at worse, threatening survival.

With recent advances in [surgical techniques](#), more patients are undergoing transcatheter [aortic valve replacement](#) (TAVR) or surgical aortic valve replacement (SAVR) to treat [aortic stenosis](#). However, this study shows that serious risks of functional decline associated with surgery in older patients may well outweigh the benefits, and in particular, may not meet an individual's goals for treatment. Although outcomes have improved over time, older patients continue to experience high rates of functional decline, particularly if they are frail before surgery. Many older patients, given the option, would often choose quality of life over longevity when making choices about their care.

Earlier [clinical trials](#) and [observational studies](#) demonstrated improvement in functional status after TAVR and SAVR. However, these studies' results may not be applicable to all patients. The studies also assessed functional limitations due to [heart failure](#) using disease-specific measures, rather than measures that looked at, for example, how patients were doing with activities of daily living, capturing a broader picture of how patients were fairing after surgery.

As this study shows, the functional status of a patient before surgery

impacts post-operative outcomes, including rates of recovery and long-range functional status. After heart valve replacement, five paths were identified that describe how much patients recover. After minimally invasive transcatheter replacement, many patients did not have improvement in functional status. In particular, patients who were very frail before their procedures tended to have the worst outcomes. Patients who are stronger generally get open heart surgery, and after surgical valve replacement, patients generally did better. Discussion of expected recovery can help to inform patient centered decision making.

According to Dr. Kim, "Although [older patients](#) with severe aortic valve stenosis undergo these procedures with expectation that their functional status improves, those with severe frailty, have functional decline. We emphasize the various processes of "how" they get to where they are at 12 months, and identified preoperative frailty, postoperative complications and delirium as strong predictors of functional decline, which can be potential targets for intervention."

Provided by Hebrew SeniorLife Institute for Aging Research

Citation: New study seeks to guide clinical treatment for older patients with aortic stenosis (2019, February 4) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-02-clinical-treatment-older-patients-aortic.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--