

Clinical trial: Prenatal DHA prevents bloodpressure increase from obesity during childhood

February 22 2019



The clinical trial's key finding was that being overweight and obese were associated with the expected higher blood pressure in the placebo group but not in the group whose mothers were assigned to DHA. Obese and overweight children of mothers in the placebo group had a large mean increase of 3.94 mm



Hg for systolic BP and 4.97 mm Hg for diastolic BP compared with overweight/obese children of DHA-supplemented mothers. These differences were statistically significant. Credit: KU / University of Kansas

Research published today in *JAMA Network Open* by investigators at the University of Kansas and KU Medical Center finds pregnant mothers who daily consumed 600 milligrams of docosahexaenoic acid (DHA)—an omega-3 fatty acid found in prenatal vitamins, fish-oil supplements and fish meat—protected their offspring from the blood pressure-elevating effects of excessive weight in early childhood.

Women with low-risk pregnancies in the Kansas City area were enrolled in the study at the University of Kansas Medical Center's Maternal and Child Nutrition and Development Lab between March 2006 and September 2009. Half were randomly assigned to a daily prenatal supplement of 600 milligrams DHA, and half were given a placebo. The primary outcomes of the intervention were pregnancy outcome and child <u>development</u> through age 6; blood pressure measured longitudinally at 4, 4 1/2, 5, 5 1/2 and 6 years of age in 171 children was a secondary outcome.

The key finding was that being overweight and obese were associated with the expected higher blood pressure in the placebo group but not in the group whose mothers were assigned to DHA. Obese and overweight children of mothers in the <u>placebo group</u> had a large mean increase of 3.94 mm Hg for systolic BP and 4.97 mm Hg for diastolic BP compared with overweight/obese children of DHA-supplemented mothers. These differences were statistically significant.

While many prenatal supplements in the U.S. contain DHA, most have much less than 600 milligrams. The researchers emphasize the amount



of prenatal DHA needed to protect against the rise in blood pressure in children with excessive weight is not known. However, being overweight and obese are large problems among U.S. children with nearly 1 in 5 school-age children and young people aged 6 to 19 years obese, according to the Centers for Disease Control 2015-2016 data.



Research published today in *JAMA Network Open* by investigators at the University of Kansas and KU Medical Center finds pregnant mothers who daily consumed 600 milligrams of docosahexaenoic acid (DHA) -- an omega-3 fatty acid found in prenatal vitamins, fish-oil supplements and fish meat -- protected their offspring from the blood pressure-elevating effects of excessive weight in early childhood. Credit: KU / University of Kansas



"This research is aimed at expectant mothers and pediatricians who wonder what you can do prior to the birth of your child to optimize health and behavior outcomes," said co-author John Colombo, KU professor of psychology, director of KU's Life Span Institute and currently KU's interim vice chancellor for research. "There's a phenomenon called 'developmental programming,' and researchers have studied effects of the prenatal environment on long-term outcomes since World War II. The prenatal environment programs a fetus' metabolism for what to expect in the postnatal environment. Part of DHA's known effects may be in programming cardiac function that preserves normal blood pressure in the case of high postnatal weight gain."

"Prenatal DHA exposure appears to program the developing fetus to be protected against the blood pressure-elevating effects of obesity in childhood," said co-author Susan Carlson, AJ Rice Professor of Nutrition in the KU Department of Dietetics and Nutrition at the KU Medical Center.

The investigators believe lower <u>blood</u> pressure at age 6 might extend beyond childhood.

"It is known that <u>blood pressure</u> tracks over time such that people with higher BP early in life are more likely to have higher BP later in life," Carlson said.





'This research is aimed at expectant mothers and pediatricians who wonder what you can do prior to the birth of your child to optimize health and behavior outcomes,' said co-author John Colombo. Credit: KU / University of Kansas

Colombo and Carlson's colleagues on the study were Elizabeth Kerling, Jamie Hilton and Jocelynn Thodosoff of the KU Department of Dietetics & Nutrition and Jo Wick of the Department of Biostatistics at KU's School of Medicine.

The National Institutes of Health supported the phase 3 clinical trial, which also is part of KU's Intellectual and Developmental Disabilities Research Center.



Previous reports of the trial found that maternal DHA supplementation reduced birth before 34 weeks' gestation and was associated with higher fat-free body mass at age 5.

More information: JAMA Network Open (2019). DOI: 10.1001/jamanetworkopen.2019.0088

Provided by University of Kansas

Citation: Clinical trial: Prenatal DHA prevents blood-pressure increase from obesity during childhood (2019, February 22) retrieved 20 May 2024 from https://medicalxpress.com/news/2019-02-clinical-trial-prenatal-dha-blood-pressure.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.