


Shedding light—literally—on resistance to radiation therapy

February 28 2019



Credit: CC0 Public Domain

A new Johns Hopkins study offers promise towards someday being able to non-invasively examine changes in cancerous tumors to determine whether they'll respond to radiation treatment, before treatment even begins.

The findings, published Feb. 28 in *Cancer Research*, can potentially help identify subsets of patients that are best suited for radiation treatment, thereby reducing medical overtreatment and helping patients avoid the adverse effects of radiation.

"Our eventual hope is to be able to predict a patient's response before [radiation therapy](#) even begins, thus sparing patients whose tumors can't be treated with radiation from going through the arduous multi-week process, saving them both time and money," says Ishan Barman, a Johns Hopkins University assistant professor of mechanical engineering and one of the study's corresponding authors.

In typical radiation therapy, small doses of radiation are delivered over a period of five to seven weeks and there's currently no definitive way of determining treatment response before or in the early weeks of therapy. The earliest clinicians can examine [tumor shrinkage](#) is two to three weeks after therapy with CT and MRI scans.

While other studies have investigated molecular changes in tumors as a way to preemptively identify tumors that won't respond to radiation treatment, the researchers say their study is unique because they studied multiple types of cancers and used smaller doses of radiation, which more accurately reflects current radiation therapy practices.

"Previous researchers used a single large dose of radiation, resulting in biomolecular changes that don't accurately mimic the subtle changes caused by smaller doses delivered over longer periods of time," says Santosh Paidi, a Johns Hopkins University mechanical engineering graduate research assistant and the study's first author.

In this study, the Johns Hopkins, University of Arkansas and the University of Arkansas for Medical Sciences researchers used Raman spectroscopy, a method which uses light from a laser to examine how molecules vibrate, to characterize the changes in the biochemical composition of the tumor and its environment. This method, the researchers say, requires only a low-power laser and doesn't need any additional sample preparation; such technologies enable holistic examination of the complex biological processes of [radiation treatment](#).

To test the different responses between tumors that are sensitive to radiation and those that are resistant to radiation, the researchers used cell lines of both resistant and sensitive tumors from human lung as well as head and neck cancers to grow tumors in mice.

After the tumors grew to a certain size, the research team removed them and scanned them with a Raman spectroscopy system. All four tumor types showed changes in response to small doses, meant to replicate clinical radiation therapy practice in humans, given over the course of two weeks.

The researchers also found consistent changes in lipid and collagen content in both lung, and head and neck tumors.

For the second part of the experiment, the research team examined untreated lung tumors. Raman spectroscopy provided data that allowed the research team to distinguish between radiation-resistant and radiation-sensitive tumors, thus providing key insight into the roles different

molecules may play in making tumors resistant to radiation.

The research team found subtle differences in each tumor type's spectrographic signature; by analyzing these patterns, the researchers created an algorithm that could identify radiation resistance and sensitivity with a 97 percent success rate.

"This is only the first step of a larger research endeavor to determine how head and neck cancer tumors respond to radiation," says Barman. "The ultimate goal is to build a miniature probe that can fit into a laryngoscope. Hopefully in the future, then, when a clinician performs an endoscopy and looks at a patient's cancerous tumor, they'll be able to determine whether that [tumor](#) will even respond to [radiation](#) therapy, and that can improve treatment plans."

Provided by Johns Hopkins University

Citation: Shedding light—literally—on resistance to radiation therapy (2019, February 28) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2019-02-lightliterallyon-resistance-therapy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--