

Psychosocial stress, the unpredictability schema, and cardiovascular disease in women

February 28 2019

In the current issue of Cardiovascular Innovations and Applications (Special Issue on Women's Cardiovascular Health, Volume 3, Number 4, 2019, Guest Editor Gladys P. Velarde) pp. 391-401(11); DOI: <https://doi.org/10.15212/CVIA.2017.0065> Tomás Cabeza de Baca, Ph.D. and Michelle A. Albert, MD, MPH from the Department of Medicine, University of California, San Francisco, CA, USA consider psychosocial stress, the unpredictability schema, and cardiovascular disease in women.

The authors provide a thorough review of the impact of psychosocial stressors stressing the importance of psychosocial factors, such as anxiety, depression, and inadequate social economic resources and their impact on adverse cardiovascular outcomes in women. This topic, often neglected, is an important contributing risk factor, especially in [ethnic minority women](#).

Depression/anxiety-related disorders and psychosocial stress have been implicated as cardiovascular disease (CVD) [risk factors](#). Women are at considerable risk for affective disorders and report greater severity from psychosocial stress, compared to men. Affective disorders and cardiovascular disease likely share underlying pathophysiological mechanisms that are potentiated among women—especially younger women. Environmental stressors that threaten the safety, security, and status of an individual are appraised by the brain, producing a cascade of evoked physiological and cognitive responses. In the short term, these processes overcome stressors, but come with long-term health

implications. Chronic [psychosocial stress](#) leads to a dysregulation of the stress response systems that can lead to a heightened stress appraisal schema called the unpredictability schema, a construct that might arguably place women at heightened risk for CVD.

This article forms part of a special issue on Women's Cardiovascular Health, guest edited by Gladys P. Velarde. Recent decades have witnessed great progress in the treatment of [cardiovascular disease](#) (CVD). Due to improved therapies, preventive strategies and increased public awareness, CVD (stroke, [heart failure](#), ischemic heart disease, peripheral arterial disease and congenital heart disease) mortality has been on the decline over this span of time for both genders.

Unfortunately, the decline has been less prominent for women, especially women of color. Once viewed as a man's disease, CVD remains the leading cause of mortality for women in the United States and is responsible for a third of all deaths of women worldwide and half of all deaths of women over 50 years of age in developing countries. In the United States, CVD far outpaces all other causes of death, including all forms of cancer combined. The statistics are sobering with about one female death in the United States every 80 seconds from CVD. That represents close to 400,000 deaths per year according to the more recent statistics. Of these, more than one quarter of a million women will die this year from [ischemic heart disease](#) (IHD) which includes obstructive and non-obstructive coronary disease, and about 64% of women who die suddenly of IHD have no prior symptoms. Despite a significant number of females with known CVD and increased awareness among women of heart disease as their major health threat, a substantial proportion of women (46% as per the most recent American Heart Association survey) remain unaware of their cardiovascular risk and continue to fail to recognize its significance.

This lack of awareness is more profound (over 60% unaware) among women in higher-risk groups, racial and ethnic minorities, and has

changed little in decades.

Poorly understood sex/gender differences in pathobiologic mechanisms, clinical presentation, management and application of diagnostic and therapeutic and preventive strategies have contributed to this gap. A critically important factor has been the underrepresentation of women in CVD research to date. In fact, only one-third of CVD clinical trials report sex-specific results despite The Food and Drug Administration regulations requiring sex stratification data, as well as the National Institute of Health recommendations of increased inclusion of women in clinical trials. This makes it difficult for researchers and clinicians to draw accurate conclusions about sex differences in mechanisms of disease, accuracy of specific diagnostic modalities and risks or benefits of a particular drug or device for the treatment of women with CVD. Furthermore, physicians and other healthcare providers continue to underestimate women's cardiovascular risk, in part because of utilization of traditional approaches which can lead to over-testing or inappropriate risk assessment without accurate differentiating who is truly at risk and inadequate use of preventive therapies for women.

The goal of this special edition Cardiovascular Innovations and Applications is to shed some light on specific topics that dominate the spectrum of CVD in [women](#).

More information: Tomás Cabeza de Baca et al, Psychosocial Stress, the Unpredictability Schema, and Cardiovascular Disease in Women, *Cardiovascular Innovations and Applications* (2019). [DOI: 10.15212/CVIA.2017.0065](https://doi.org/10.15212/CVIA.2017.0065)

Provided by Cardiovascular Innovations and Applications

Citation: Psychosocial stress, the unpredictability schema, and cardiovascular disease in women (2019, February 28) retrieved 20 April 2024 from <https://medicalxpress.com/news/2019-02-psychosocial-stress-unpredictability-schema-cardiovascular.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.