


Study finds low statin use among some high-risk kidney disease patients

February 19 2019



Credit: CC0 Public Domain

Clinical trials have shown that cholesterol-lowering statin drugs can significantly reduce the risk of cardiovascular disease in kidney disease patients who are not on dialysis.

But a new study has found that statins are used by only 21.8 percent of

such patients who do not already have [cardiovascular disease](#) or diabetes or have not been diagnosed with high cholesterol.

The study by researchers from Loyola Medicine, Loyola University Chicago and Hines VA Hospital was published in *Clinical Kidney Journal*.

"Our findings suggest a need for education efforts to increase statin use in adults with non dialysis-dependent chronic kidney disease," reported first author Talar Markossian, Ph.D., MPH, corresponding author Holly Kramer, MD, MPH, and colleagues.

More than 500 million people worldwide have [chronic kidney disease](#) and are not on dialysis. Most cases are due to diabetes and [high blood pressure](#), which also are strong risk factors for heart attacks, strokes and other forms of cardiovascular disease. The risk of cardiovascular disease increases sharply as kidney disease becomes worse.

In the new study, researchers examined records of 581,344 patients receiving care from Veterans Affairs facilities. Patients were aged 50 and older, with stages 3-5 kidney disease that did require dialysis.

About three-fourths of patients who had kidney disease plus cardiovascular disease or diabetes used statins. But in the absence of cardiovascular disease, diabetes or high cholesterol, statin use was only 21.8 percent. This finding is concerning because about 90 percent of such patients have at least a 10 percent risk of developing cardiovascular disease in the next 10 years, and this risk could be reduced by taking statins. Statin use also is believed to be low in patients outside the VA.

More information: Talar W Markossian et al, Low statin use in nondialysis-dependent chronic kidney disease in the absence of clinical atherosclerotic cardiovascular disease or diabetes, *Clinical Kidney*

Journal (2019). [DOI: 10.1093/ckj/sfy131](https://doi.org/10.1093/ckj/sfy131)

Provided by Loyola University Health System

Citation: Study finds low statin use among some high-risk kidney disease patients (2019, February 19) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-02-statin-high-risk-kidney-disease-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.