

Team improves emergency care for patients with chest pain

February 27 2019

Emergency physicians at Kaiser Permanente hospitals in Southern California reduced hospital admissions and cardiac stress testing by using new criteria to assess the level of risk patients with chest pain have for subsequent cardiac events.

The [emergency department](#) assessment was implemented across Kaiser Permanente Southern California and has improved the quality and efficiency of care for these patients according to an article published today in *Annals of Emergency Medicine*.

"By assessing patients who come into the emergency department using HEART—a measure of history, electrocardiogram (ECG), age, [risk factors](#) and troponin, a [blood test](#) that measures heart injury—we were able to avoid unnecessary hospitalizations and cardiac testing for many patients without increasing their risk of heart damage," said Adam L. Sharp, MD, MSc, an emergency physician at the Kaiser Permanente Los Angeles Medical Center and a researcher with the Kaiser Permanente's Department of Research & Evaluation in Southern California.

Heart disease is a leading cause of death in the United States. Chest pain—a symptom that often triggers an evaluation for suspected acute coronary syndrome, or [heart attack](#)—is the second most common reason for emergency department visits. These patients are frequently hospitalized for observation and cardiac stress testing, but no studies have evaluated whether this strategy improves patient health. Researchers involved in this study sought to evaluate whether the

HEART tool could successfully identify which patients required observation and further testing, and which ones could be allowed to safely go home.

Researchers used information in the electronic health records of more than 65,000 adult patients evaluated for suspected [acute coronary syndrome](#) at 13 Kaiser Permanente emergency departments in Southern California between May 2015 and June 2017.

The study examined outcomes in 3 periods:

- May 2015 to May 2016 before HEART was established
- May 6, 2016 to June 3, 2016, during the transition to HEART
- June 2016 to June 2017 after HEART was implemented

The outcomes studied were hospital admissions, non-invasive cardiac stress testing, and whether patients died or had heart attacks within 30 days of the emergency department visit.

Researchers found that, among patients who went to the emergency department with chest pain, there was:

- A 4.4 percent decrease in hospital admissions and cardiac stress testing after implementation of HEART.
- No increase in missed heart attacks in patients after HEART was implemented, which was already below the nationally reported missed rate.
- No increase in the rate of death in patients after HEART was implemented.

"Our program to decrease hospitalization of patients with low-risk [chest pain](#) builds upon previous efforts to improve care by avoiding unwarranted, potentially harmful treatments such as reducing the

prescriptions of antibiotics for acute sinusitis and reducing diagnostic tests such as CT imaging after head trauma," Dr. Sharp said.

Provided by Kaiser Permanente

Citation: Team improves emergency care for patients with chest pain (2019, February 27)
retrieved 6 May 2024 from

<https://medicalxpress.com/news/2019-02-team-emergency-patients-chest-pain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.