


In vitro grafts increase blood flow in infarcted rat hearts

February 5 2019, by Brian Donohue



Histology slides show murine heart at five days post-implant a) with conventional non-perfusible vessels and b) with patterned perfusable cardiac construct. Credit: Nicole Zeinstra / University of Washington Bioengineering

Advances in stem cell research offer hope for treatments that could help

patients regrow heart muscle tissue after heart attacks, a key to achieving more complete recovery.

Scientists today report success in creating functional [blood vessels](#) in vitro for hearts of rats that had sustained a heart attack. The journal *Nature Communications* published the paper, whose lead authors are Ying Zheng and Charles Murry of the UW Medicine Institute for Stem Cell and Regenerative Medicine in Seattle.

"To our knowledge, this is the first demonstration that building organized blood vessels with perfusion outside the body leads to improved integration with host blood vessels and better [tissue](#) blood flow," said Zheng, a University of Washington associate professor of bioengineering.

The scientists set out to show that by growing stem cell-derived heart tissue in a petri dish, with attention to blood vessels' construction, they could improve the tissue's incorporation with existing heart vessels.

"I come from a mechanical background," Zheng continued. "I love thinking about the dynamics of blood flow. Our whole bodies are vascularized. This network of vessels is dynamic and interconnected, like a transportation system that remodels itself all the time."

Disruption to blood flow during a [heart attack](#) leads to significant loss of heart muscle and heart function. Heart muscle grown from stem cells must not only survive and integrate with the host tissue, but it must also restore adequate blood flow, explained Murry. He is a UW professor of pathology, bioengineering, and medicine/cardiology.

The research team used human [stem cells](#) to create a vascularized construct, or patch, with a functioning network of blood vessels that mimics the vasculature of a human heart.

"Being able to organize the vessels in the tissue outside the body was very important," Zheng said. "When we implanted the patch, we saw that the stem cell-derived tissue integrated effectively with the host's coronary circulation. This improved blood flow to the engineered tissue and gave it the nutrients it needed to survive."

Optical microangiography imaging techniques developed by Ricky Wang, a UW professor of bioengineering, revealed that blood flow within the grafts was twentyfold higher than has been reported for any other such graft. This suggested that nurturing the tissue in the lab had a meaningful benefit for the [heart](#) cells before they were implanted into the rats' hearts, the researchers said.

More information: Meredith A. Redd et al, Patterned human microvascular grafts enable rapid vascularization and increase perfusion in infarcted rat hearts, *Nature Communications* (2019). [DOI: 10.1038/s41467-019-08388-7](https://doi.org/10.1038/s41467-019-08388-7)

Provided by University of Washington

Citation: In vitro grafts increase blood flow in infarcted rat hearts (2019, February 5) retrieved 23 April 2024 from <https://medicalxpress.com/news/2019-02-vitro-grafts-blood-infarcted-rat.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--