

Blood pressure control reduces dementia risk in mid-life patients with atrial fibrillation

March 18 2019

Dementia risk in mid-life patients with atrial fibrillation (AF) can be reduced by controlling high blood pressure, according to a study presented today at EHRA 2019, a congress of the European Society of Cardiology (ESC).

Study author Dr. Daehoon Kim said that AF patients in their 50s and 60s can benefit from lowering their [systolic blood pressure](#) (SBP) to 120-129 mmHg, as compared to those with continuously high SBP over 140mmHg.

"It was surprising. We all know that AF is associated with higher risk of [dementia](#), but there was no certain way to protect against AF-associated dementia," explained Dr. Kim, a cardiologist at Yonsei University College of Medicine in Seoul, Republic of Korea. "With this study, we found one potential way to help mid-life patients minimise their risk by as much as 15%," he said.

AF patients with hypertension and taking at least one antihypertensive medication at the time of AF diagnosis between 2005-2016 were enrolled in the Korean nationwide cohort study. Among those with uncontrolled SBP over 140mmHg at initial AF diagnosis, the risk of dementia was compared according to their follow-up SBP control status (

Citation: Blood pressure control reduces dementia risk in mid-life patients with atrial fibrillation

(2019, March 18) retrieved 18 April 2024 from <https://medicalxpress.com/news/2019-03-blood-pressure-dementia-mid-life-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.