

Chest CT may be better than PET/CT for HNSCC distant metastasis evaluation

March 11 2019



(HealthDay)—As distant metastasis (DM) in head and neck squamous

cell carcinoma (HNSCC) is a rare event, computed tomography of the chest may be more cost-effective for evaluation, according to a study recently published in *Oral Oncology*.

Jeffrey C. Liu, M.D., from Temple University in Philadelphia, and colleagues examined the frequency and location of DM at presentation in HNSCC using data from the National Cancer Data Base (2010 to 2015) for 151,730 cases.

The researchers found that the nasopharynx (NP) had the highest percentage of M1 disease (9.1 percent), followed by the hypopharynx (7.3 percent). M1 cases were reported in 3.1 percent of the study population when excluding the NP. For all subsites ([oral cavity](#), oropharynx, larynx, hypopharynx, and nasopharynx), advanced T-stage, positive N-stage, and N3 status were all predictors of M1 status. The majority of DM cases (53.1 percent) presented with at least lung involvement when metastasis site was known. However, in NP cancers, just under one-third (32.8 percent) of DM included the lung.

"Our findings confirm that checking for distant metastasis in this group of patients using a chest computed [tomography](#) instead of a [positron emission tomography](#)/computed tomography scan would save patients time and money, and spare them from a more involved procedure," a coauthor said in a statement.

More information: [Abstract/Full Text](#)

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Chest CT may be better than PET/CT for HNSCC distant metastasis evaluation (2019, March 11) retrieved 3 May 2024 from <https://medicalxpress.com/news/2019-03-chest-ct-petct-hnscc-distant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.