

Common chest infection puts babies at risk of hospitalization for asthma in preschool years

March 5 2019

Infants who are admitted to hospital with the common infection bronchiolitis are at increased risk of further emergency hospital admissions for asthma, wheezing and respiratory illness in the first five years of their life.

These are the findings of researchers from Imperial College London, who tracked 613,377 babies (almost all births in England between April 2007—March 2008) up to the age of five years. Around 16, 000 babies were admitted to hospital with [bronchiolitis](#) before their first birthday.

Researchers found these children had up to a fivefold risk of emergency admissions to hospital with [asthma](#), wheezing or a respiratory illness in the first five years of their life, compared with those who were not admitted for bronchiolitis.

The researchers found that one in five children admitted to hospital with bronchiolitis as a baby will have a subsequent emergency hospital [admission](#) for asthma, wheeze, or respiratory infections.

Dr. Helen Skirrow, lead author of the research from the School of Public Health at Imperial, said: "We know that hospital admissions for bronchiolitis have been rising over recent years in England. Previous studies have suggested a link between the condition and an increased risk of conditions such as asthma, but this is the largest study to suggest a

severe case of bronchiolitis can result in subsequent emergency admissions for asthma and other respiratory conditions. "

Dr. Skirrow added: "An emergency hospital admission is incredibly stressful for children and their families, and also places a burden on hospitals. If we develop interventions to prevent the initial bout of bronchiolitis—we may also be able to reduce the number of subsequent emergency admissions."

Bronchiolitis is a common type of chest infection that affects around one in three children in their first year of life, most commonly babies between three and six months of age.

The condition causes the airways to become inflamed, triggering symptoms such as a fever, cough and wheezing. The majority of children get better on their own—however up to one in 20 (2-5%) will require hospital treatment if they develop more severe symptoms, such as breathing difficulties.

Most cases of bronchiolitis are caused by the respiratory syncytial virus (RSV) which is responsible for around 3.2 million hospital admissions every year worldwide. Although there is no treatment for bronchiolitis, children thought to be at risk of RSV infection, such as babies born prematurely, can be given a preventative treatment called palivizumab. There are also a number of RSV vaccines currently in development.

Many factors contribute to a baby's risk of being admitted to hospital for bronchiolitis and other [respiratory illnesses](#). These include exposure to tobacco smoke inside the home, environmental factors (such as air quality) and whether a child was breastfed.

The team say this current finding, published in the journal *Archives of Disease in Childhood*, highlight the importance of preventing cases of

bronchiolitis.

Professor Sonia Saxena, senior author and head of Imperial's Child Health Unit, added: "Having your baby admitted to hospital with a serious chest infection like bronchiolitis is a frightening experience for parents. Our study shows these children are at high risk of subsequent [hospital](#) admission for asthma and recurrent infections across the early years of life, so we should be doing all we can to prevent this happening. Health professionals should counsel parents about benefits of breastfeeding, avoid infant's exposure to [tobacco smoke](#) and raise parent's awareness of when and how to seek help for respiratory problems as their child develops during preschool years."

When analysing the findings the research team took into account infants with pre-existing conditions. The scientists say more work is still needed to investigate why bronchiolitis may be linked to subsequent respiratory problems such as asthma. One theory is the initial viral infection that caused bronchiolitis may alter the immune system's response, increasing the chance of subsequent asthma and wheezing. Or it may be that a baby remains exposed to the initial triggers of bronchiolitis, such as second-hand smoke or pollution, which subsequently triggers asthma or other respiratory conditions.

The team say the research also highlights the need to develop a vaccine against RSV, which would help prevent cases of bronchiolitis.

Dr. Skirrow explained: "There are a number of trials underway at the moment developing an RSV vaccine that could either be given to newborns or pregnant women. This work provides more evidence of the importance of investing in this research, and the effect it could have on children's long-term health."

Provided by Imperial College London

Citation: Common chest infection puts babies at risk of hospitalization for asthma in preschool years (2019, March 5) retrieved 12 May 2024 from

<https://medicalxpress.com/news/2019-03-common-chest-infection-babies-hospitalization.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.