

Engaging in physical activity could reduce long-term mortality

March 12 2019



Irene Esteban Cornejo. Credit: University of Granada

Cognitive frailty is a heterogeneous clinical manifestation characterized by the simultaneous presence of both physical frailty and cognitive impairment, in the absence of dementia, and it seems to entail a greater death risk than physical frailty or cognitive impairment separately. Despite the potential effect of regular physical activity to slow cognitive

decline and its association with lower mortality in nonfrail individuals, no previous studies have investigated whether and to what extent physical activity could attenuate the effect of cognitive frailty on mortality.

Colleagues from the University of Granada, Universidad Autónoma de Madrid, IMDEA Food Institute, CIBER in Epidemiology and Public Health and Northeastern provide the first empirical evidence about the impact of [physical activity](#) on mortality in cognitively frail individuals. They studied 3677 older adults with a mean age of 72 years, who were followed up for 14 years. These novel results showed that cognitive frailty was associated with increased mortality more markedly in the inactive older adults and, more importantly, being physically active may reduce the mortality risk among cognitively frail individuals by 36 percent.

Irene Esteban-Cornejo, first author of the study and researcher from the University of Granada highlights the clinical implications of the study and notes that: "Compared to those who were robust and active, participants with cognitive frailty who were inactive had the highest mortality risk, which was equivalent to being almost 7 years older."

This research may have important implications because the levels of physical activity are dramatically reduced in the elderly while cognitive and physical functioning decline is naturally occurring, stated David Martinez-Gómez, last author of the study and researcher from the Autonomous University of Madrid. In the present study, the prevalence of physical inactivity in cognitively frail individuals was around 65 percent. Therefore, promoting physical activity among cognitively frail elderly persons is crucial because they have room for improvement and it may increase their survival. However, further longitudinal and experimental studies are needed to shed light on the importance of physical activity to decrease mortality in cognitively frail individuals.

Esteban-Cornejo and colleagues conclude that promoting a physically [active lifestyle](#) could be one of the main strategies to improve vital prognosis among cognitively frail [older adults](#). They call to health Institutions to prescribe physical activity as a treatment against cognitive frailty-related [mortality](#).

More information: Irene Esteban-Cornejo et al. Cognitive Frailty and Mortality in a National Cohort of Older Adults: the Role of Physical Activity, *Mayo Clinic Proceedings* (2019). [DOI: 10.1016/j.mayocp.2018.10.027](#)

Provided by University of Granada

Citation: Engaging in physical activity could reduce long-term mortality (2019, March 12) retrieved 6 May 2024 from <https://medicalxpress.com/news/2019-03-engaging-physical-long-term-mortality.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--