

## Excess hormones could cause a condition that can lead to blindness in women, study finds

March 21 2019



Credit: CC0 Public Domain

Research led by the University of Birmingham has found that increased levels of hormones including testosterone could cause a brain condition that can lead to blindness in women.

Idiopathic Intracranial Hypertension—also known as IIH—is caused by <u>high pressure</u> in the brain with consequences from blindness to



incapacitating daily long-term headaches. IIH was originally identified over 100 years ago yet the cause of the condition has remained unknown although there has been much speculation about why more than 95 per cent of total incidence is in <u>women</u> with obesity.

Once regarded as rare, incidence of IIH is now rising dramatically in line with the global rise in obesity and there has been a 150 per cent rise in incidence in last 10 years. Currently there are very limited treatment options.

Scientists at the University of Birmingham carried out research, funded by the National Institute of Health Research (NIHR), the Medical Research Council UK, and Wellcome Trust and published today in *JCI Insight*, exploring the role of hormones known as androgens in the way IIH develops.

The researchers examined the levels of androgens in blood and urine, as well as in brain fluid known as cerebrospinal fluid (CSF), in 55 women with IIH aged 18 to 45. They then compared the results with the levels observed in women with obesity of the same age and body mass index (BMI), as well as a cohort of women with polycystic ovary syndrome (PCOS).

PCOS is a common condition and is classically associated with increased levels of specific types of androgens, which can cause excess hair and irregular periods.

The researchers found that IIH patients had a very different profile of raised levels of androgens compared to women of a similar age with obesity and women with PCOS. Most notable were the high levels of the androgen 'testosterone' found in the blood in IIH women. Crucially, levels of androgens were uniquely increased in the brain fluid (CSF) of women with IIH. When the researchers, analysed human choroidal



plexus tissue, which is the site in the brain where CSF is produced, they confirmed that androgens could increase the rate of CSF secretion, a potential driver for increased brain pressure.

Corresponding and senior author Dr. Alex Sinclair, NIHR Clinician Scientist at the University of Birmingham, said: "For over 100 years the cause of IIH has been unknown.

"These findings mark a key step forward; for the first time we have found a pattern of androgen dysregulation that is unique to IIH and potentially a driver of abnormal brain pressure in people with the condition.

"Taken together this provides the first evidence that IIH may be a manifestation of female <u>androgen</u> excess."

Lead author Dr. Michael O'Reilly, Hoffenberg Clinician Scientist in Endocrinology and Human Metabolism at the University of Birmingham, said: "These findings, for the first time, implicate androgens as a potential cause of IIH."

Professor Wiebke Arlt, Director of the University of Birmingham's Institute of Metabolism and Systems Research, added: "These findings mark a significant step forward in a previously poorly understood disabling condition.

"Understanding the cause of a disease is a vital step to ultimately refine treatments which improved patient care and outcomes."

Michelle Williamson, Chair of the patient charity IIHUK, added: "The recent patient and doctor research Priority Setting Partnership identified that uncovering the cause of IIH was the number one priority for IIH. We are delighted by this discovery. It is vital for the patients to see that



progress is being made."

This latest study comes after experts led by the University of Birmingham recently produced the first internationally collaborative guidelines aimed at improving the treatment and diagnosis of IIH.

Created by a group of specialists in neurology, neurosurgery, neuroradiology, ophthalmology and neuroradiology, who have had expertise in managing IIH, the team envisage that the new guidelines, published last October in *Journal of Neurology Neurosurgery and Psychiatry*, will step change the treatment of the condition.

Provided by University of Birmingham

Citation: Excess hormones could cause a condition that can lead to blindness in women, study finds (2019, March 21) retrieved 3 May 2024 from https://medicalxpress.com/news/2019-03-excess-hormones-condition-women.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.